

## A SEMILINEAR FOURTH ORDER ELLIPTIC PROBLEM WITH EXPONENTIAL NONLINEARITY\*

GIANNI ARIOLI<sup>†</sup>, FILIPPO GAZZOLA<sup>†</sup>, HANS-CHRISTOPH GRUNAU<sup>‡</sup>, AND  
ENZO MITIDIERI<sup>§</sup>

**Abstract.** We study a semilinear fourth order elliptic problem with exponential nonlinearity. Motivated by a question raised in [P.-L. Lions, *SIAM Rev.*, 24 (1982), pp. 441–467], we partially extend results known for the corresponding second order problem. Several new difficulties arise and many problems still remain to be solved. We list those of particular interest in the final section.

**Key words.** biharmonic operator, super-solutions, computer assisted proof

**AMS subject classifications.** 35G30, 35J40

**DOI.** 10.1137/S0036141002418534

**1. Introduction.** In the last forty years a great deal has been written about existence and multiplicity of solutions to nonlinear second order elliptic problems in bounded and unbounded domains of  $\mathbb{R}^n$  ( $n \geq 2$ ). Important achievements on this topic have been made by applying various combinations of analytical techniques, and among all of them we mention only the variational and topological methods. For the latter, especially when the main interest is focused on the existence of positive solutions, the fundamental tool which has been used is the maximum principle [A1] and its consequences [GNN].

For higher order problems, a possible failure of the maximum principle causes several technical difficulties. This fact is very likely the reason why the knowledge on higher order nonlinear problems is far from being reasonably complete, as it is in the second order case.

One of the most interesting and intensively studied second order model problems that exhibits several peculiar features of most nonlinear elliptic equations is the so-called Gel'fand problem [G, section 15],

$$(1) \quad \begin{cases} -\Delta u = \lambda e^u & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

Here  $\Omega$  is a bounded smooth domain in  $\mathbb{R}^n$  ( $n \geq 3$ ) and  $\lambda \geq 0$  is a parameter. This problem appears in connection with combustion theory [G, JL] and stellar structure [C]. From a mathematical point of view, one of the main interests is that it may have both unbounded (singular) solutions and bounded (regular) solutions (see [BV, GMP, MP1, MP2]): by the results in [CR, BCMR] it is known that there exists  $\lambda^* > 0$  such that if  $\lambda > \lambda^*$  there exists no solution of (1) (neither regular nor singular),

---

\*Received by the editors November 25, 2002; accepted for publication (in revised form) April 23, 2004; published electronically February 3, 2005.

<http://www.siam.org/journals/sima/36-4/41853.html>

<sup>†</sup>Dipartimento di Matematica del Politecnico, via Bonardi 9, 20133 Milano, Italy (gianni@mate.polimi.it, gazzola@mate.polimi.it). The work of these authors was supported by the MURST project “Metodi Variazionali ed Equazioni Differenziali non Lineari.”

<sup>‡</sup>Fakultät für Mathematik, Otto-von-Guericke-Universität, Postfach 4120, 39016 Magdeburg, Germany (Hans-Christoph.Grunau@mathematik.uni-magdeburg.de).

<sup>§</sup>Dipartimento di Scienze Matematiche, via A. Valerio 12/1, 34100 Trieste, Italy (mitidier@univ.trieste.it).

while if  $0 \leq \lambda < \lambda^*$ , there exists a minimal regular solution  $U_\lambda$  of (1) and the map  $\lambda \mapsto U_\lambda$  is smooth and increasing. In the unit ball  $B$ , the bifurcation picture of radial solutions is rather complete. There is always a singular solution  $u_\sigma := -2 \log |x|$  with corresponding parameter  $\lambda_\sigma = 2(n - 2)$ . If  $n \geq 10$ , the solution branch consists only of minimal solutions and terminates at  $\lambda^* = \lambda_\sigma$  in the singular solution. If  $3 \leq n \leq 9$ , then  $\lambda^* > \lambda_\sigma$  and the extremal point  $(\lambda^*, U_*)$  is a turning point. The branch bends back and meanders infinitely many times around  $\lambda_\sigma$  while approaching the singular solution  $u_\sigma$ . We refer to [BV, Figure 1] for the pictures. The interested reader may see also [BE] for an account on motivations and related results.

Some interesting generalizations of (1) have been considered in the framework of second order quasi-linear operators. We refer to [GPP] for equations associated to the  $p$ -Laplace operator and to [J, JS] for the case of the  $k$ -Hessian operator.

The aim of this paper is to give a contribution to the solution of a special case of a problem formulated in [Li, section 4.2 (c)], namely, *Is it possible to obtain a description of the solution set for higher order semilinear equations associated to exponential nonlinearities?*

Recently, interest in higher order nonlinear problems due to its exciting and promising developments has become increasingly evident especially for fourth order equations [PT]. Following this trend, we shall consider in this paper the fourth order version of (1), a semilinear elliptic problem which involves the biharmonic operator, more precisely,

$$(P_\lambda) \quad \begin{cases} \Delta^2 u = \lambda e^u & \text{in } B, \\ u = \frac{\partial u}{\partial \mathbf{n}} = 0 & \text{on } \partial B. \end{cases}$$

Here  $B$  denotes the unit ball in  $\mathbb{R}^n$  ( $n \geq 5$ ) centered at the origin and  $\frac{\partial}{\partial \mathbf{n}}$  the differentiation with respect to the exterior unit normal, i.e., in radial direction;  $\lambda \geq 0$  is a parameter. We are interested in two kinds of solutions of  $(P_\lambda)$ , regular solutions and singular solutions; see Definition 1 in the next section. We restrict our attention to the case  $n \geq 5$ , where the nonlinearity is supercritical. In low dimensions  $1 \leq n \leq 4$  the problem is subcritical and has a different behavior; see Remark 14 at the end of the following section.

Many techniques, familiar from second order equations like the maximum principle, are not available here. But since we restrict ourselves to the ball, at least a comparison principle is available; see Lemma 16 below. Moreover, in fourth order equations, one usually does not succeed in finding suitable nontrivial auxiliary functions satisfying again a differential inequality. This is a serious difficulty in proving Theorem 3 (cf. the proof of [BCMR, Theorem 3]), and it is overcome by carefully exploiting the properties of the exponential nonlinearity and the construction of minimal solutions, based upon the already mentioned comparison principle. Finally, when looking for radial solutions, one may perform a phase space analysis for the corresponding system of ODEs. Here, the phase space is no longer two-dimensional, where the topology is relatively simple and the Poincaré–Bendixson theory is available, but we have to work in a four-dimensional phase space. Some of the resulting difficulties could be overcome only with computer assistance.

This paper is organized as follows: In the next section we state some definitions and the main results contained in this work (see Theorems 3, 4, 6, 7, and 12 below). The content of sections 3 through 7 is devoted to the proofs of these theorems. Section 8 contains some results on the stability of regular solutions of  $(P_\lambda)$  and a list

of open problems that we consider of some interest and related to the main results of this paper. Finally, in section 9 we describe the algorithm used in the *computer assisted proof* of Theorem 7.

**2. Main results.** We first make precise in which sense we intend a function to solve  $(P_\lambda)$ . For this purpose, we fix some exponent  $p$  with  $p > \frac{n}{4}$  and  $p \geq 2$ . The definitions and results below do not depend on the special choice of  $p$ .

DEFINITION 1. *We say that  $u \in L^2(B)$  is a solution of  $(P_\lambda)$  if  $e^u \in L^1(B)$  and*

$$(2) \quad \int_B u \Delta^2 v = \lambda \int_B e^u v \quad \text{for all } v \in W^{4,p} \cap H_0^2(B).$$

*We say that a solution  $u$  of  $(P_\lambda)$  is regular (resp., singular) if  $u \in L^\infty(B)$  (resp.,  $u \notin L^\infty(B)$ ).*

Clearly, according to this definition, regular and singular solutions exhaust all possible solutions. Note that by standard regularity theory for the biharmonic operator (see [ADN]), any regular solution  $u$  of  $(P_\lambda)$  satisfies  $u \in C^\infty(\bar{B})$ . Note also that by the positivity preserving property of  $\Delta^2$  in the ball  $[B]$  any solution of  $(P_\lambda)$  is positive; see also Lemmas 16 and 18 below for a generalized statement. This property is known to fail in general domains. For this reason, we restrict ourselves to balls also in Theorems 3 and 4; cf. also Open Problem 8 in section 8.

We also need the notion of minimal solution, as follows.

DEFINITION 2. *We call a solution  $U_\lambda$  of  $(P_\lambda)$  minimal if  $U_\lambda \leq u_\lambda$  a.e. in  $B$  for any further solution  $u_\lambda$  of  $(P_\lambda)$ .*

In order to state our results, we denote by  $\lambda_1 > 0$  the first eigenvalue for the biharmonic operator with Dirichlet boundary conditions

$$(3) \quad \begin{cases} \Delta^2 u = \lambda_1 u & \text{in } B, \\ u = \frac{\partial u}{\partial \mathbf{n}} = 0 & \text{on } \partial B; \end{cases}$$

it is known from the mentioned positivity preserving property and Jentzsch's (or Krein–Rutman's) theorem that  $\lambda_1$  is isolated and simple and that the corresponding eigenfunctions do not change sign.

We may now state the following theorem.

THEOREM 3. *There exists*

$$\lambda^* \in \left[ 14.72(n-1)(n-3), \frac{\lambda_1}{e} \right)$$

*such that the following hold:*

(i)  $(P_\lambda)$  admits a minimal regular solution  $U_\lambda$  for all  $\lambda < \lambda^*$  and no solutions if  $\lambda > \lambda^*$ .

(ii) The map  $\lambda \mapsto U_\lambda(x)$  is strictly increasing for all  $x \in B$ . Moreover, there exists a solution  $U_*$  of  $(P_{\lambda^*})$  which is the pointwise limit of  $U_\lambda$  as  $\lambda \uparrow \lambda^*$ .

(iii)  $U_\lambda \rightarrow U_*$  in the norm topology of  $H_0^2(B)$  as  $\lambda \uparrow \lambda^*$ .

(iv) The extremal solution  $U_*$  and all the minimal solutions  $U_\lambda$  (for  $\lambda < \lambda^*$ ) are radially symmetric and radially decreasing.

It is remarkable that at  $\lambda^*$  there is an immediate switch from existence of regular minimal solutions to nonexistence of any (even singular) solution. The only possibly singular minimal solution corresponds to  $\lambda = \lambda^*$ . This result is known from [BCMR] for the second order problem (1), but the *method* used there may not be carried over

to fourth order problems. Nevertheless, the *result* extends to the biharmonic case. The proof is given in Lemma 20 below.

We may also characterize the uniform convergence to 0 of  $U_\lambda$  as  $\lambda \rightarrow 0$  by giving the precise rate of its extinction.

THEOREM 4. *For all  $\lambda \in (0, \lambda^*)$  let  $U_\lambda$  be the minimal solution of  $(P_\lambda)$  and let*

$$V_\lambda(x) = \frac{\lambda}{8n(n+2)} [1 - |x|^2]^2.$$

*Then  $U_\lambda(x) > V_\lambda(x)$  for all  $\lambda < \lambda^*$  and all  $|x| < 1$ , and*

$$\lim_{\lambda \rightarrow 0} \frac{U_\lambda(x)}{V_\lambda(x)} = 1 \quad \text{uniformly with respect to } x \in B.$$

A complete result in the spirit of Gidas, Ni, and Nirenberg [GNN] does not hold for fourth order equations under Dirichlet boundary conditions. It has been recently proved by Sweers in [Sw] that for general semilinear autonomous biharmonic equations in a ball under Dirichlet boundary conditions, we may have positive radially symmetric solutions which are not radially decreasing, provided the right-hand side is *not* positive everywhere. This phenomenon may not occur in our situation; however, it is not known whether any smooth solution of  $(P_\lambda)$  is radially symmetric. Moreover, also in the second order case it is not known whether singular solutions are always radially symmetric. Nevertheless, Theorem 3 suggests that we pay particular attention to radially symmetric solutions. In this context, we put  $r = |x|$  and consider the functions  $u = u(r)$ .

First of all, in the following definition we introduce a new notion of solution which seems to be the natural framework for radially symmetric solutions.

DEFINITION 5. *We say that a radial singular solution  $u = u(r)$  of  $(P_\lambda)$  is weakly singular if the limit  $\lim_{r \rightarrow 0} ru'(r)$  exists.*

We do not know whether every singular solution is also weakly singular. In the second order case, Joseph and Lundgren [JL] reduce (1) to a system of two ODEs and study its phase portrait in  $\mathbb{R}^2$ ; using Bendixson’s theorem, they show that singular solutions are also weakly singular. For the fourth order equation  $(P_\lambda)$  a similar argument should be carried out in  $\mathbb{R}^4$  (see section 3) where a general result of Bendixson’s type does not hold. Therefore, the equivalence between singular and weakly singular solutions seems out of reach in our context; see Open Problem 5 in section 8.

If we seek radially symmetric solutions, we rewrite problem  $(P_\lambda)$  as  $(0 < r \leq 1)$

$$(4) \quad \begin{cases} \frac{d^4u}{dr^4} + \frac{2(n-1)}{r} \frac{d^3u}{dr^3} + \frac{(n-1)(n-3)}{r^2} \frac{d^2u}{dr^2} - \frac{(n-1)(n-3)}{r^3} \frac{du}{dr} = \lambda e^{u(r)}, \\ u(1) = 0, \\ \frac{du}{dr} \Big|_{r=1} = 0. \end{cases}$$

In [GPP, JL, MP2] the second order equation (1) was reduced to a system of two autonomous ODEs. Here, we reduce (4) to a system of four equations. First, we make the change of variables

$$(5) \quad s = \log r, \quad v(s) = u(e^s), \quad s \in (-\infty, 0]$$

so that (4) becomes

$$(6) \quad \begin{cases} \frac{d^4v}{ds^4} + 2(n-4)\frac{d^3v}{ds^3} + (n^2 - 10n + 20)\frac{d^2v}{ds^2} - 2(n-2)(n-4)\frac{dv}{ds} = \lambda e^{4s+v(s)}, \\ v(0) = 0, \\ \frac{dv}{ds} \Big|_{s=0} = 0; \end{cases}$$

then we set

$$(7) \quad \begin{cases} v_1(s) = v'(s) + 4, \\ v_2(s) = -v''(s) - (n-2)v'(s), \\ v_3(s) = -v'''(s) + (4-n)v''(s) + 2(n-2)v'(s), \\ v_4(s) = -\lambda e^{v(s)+4s}. \end{cases}$$

Finally, we obtain the following (nonlinear) differential system:

$$(8) \quad \begin{cases} v_1'(s) = (2-n)v_1(s) - v_2(s) + 4(n-2), \\ v_2'(s) = 2v_2(s) + v_3(s), \\ v_3'(s) = (4-n)v_3(s) + v_4(s), \\ v_4'(s) = v_1(s)v_4(s) \end{cases}$$

with initial conditions

$$(9) \quad v_1(0) = 4, \quad v_4(0) = -\lambda.$$

It turns out that (8) admits only the two stationary points  $P_1 = (4, 0, 0, 0)$  and  $P_2 = (0, 4n - 8, 16 - 8n, -8(n - 2)(n - 4))$ ; see section 3.1. Then, in section 3.2, we prove the following result.

**THEOREM 6.** *Let  $u = u(r)$  be a radial solution of  $(P_\lambda)$  and let*

$$V(s) = (v_1(s), v_2(s), v_3(s), v_4(s))$$

*be the corresponding trajectory relative to (8). Then*

(i)  *$u$  is regular (i.e.  $u \in L^\infty(B)$ ) if and only if*

$$\lim_{s \rightarrow -\infty} V(s) = P_1;$$

(ii)  *$u$  is weakly singular if and only if*

$$\lim_{s \rightarrow -\infty} V(s) = P_2.$$

Our following results concern the *existence* of weakly singular solutions and a lower bound  $\lambda_{\min}^*$  on the value of  $\lambda^*$ . For all  $n = 5, \dots, 16$  we prove the existence of  $\lambda_\sigma$  such that  $(P_{\lambda_\sigma})$  admits a weakly singular solution; we provide a lower and upper bound on the value of  $\lambda_\sigma$ . For all  $n = 5, \dots, 16$  let  $\lambda_\sigma^{\min}$  and  $\lambda_\sigma^{\max}$  be as given in Table 1, and for all  $n = 5, \dots, 10$  let  $\lambda_{\min}^*$  be as given in Table 1.

**THEOREM 7.** *For all  $n = 5, \dots, 16$  there exists  $\lambda_\sigma \in [\lambda_\sigma^{\min}, \lambda_\sigma^{\max}]$  such that  $(P_{\lambda_\sigma})$  admits a weakly singular solution  $U_\sigma$ . In particular,  $\lambda_\sigma > 8(n - 2)(n - 4)$ .*

*For all  $n = 5, \dots, 10$  the value of  $\lambda^*$  is larger than  $\lambda_{\min}^*$ .*

In section 6 we use Theorem 6 to show that Theorem 7 is equivalent to some intersection properties of the unstable manifolds of  $P_1$  and  $P_2$  with the hyperplane

TABLE 1

$n$	$\lambda_\sigma$	$\lambda^*$	$\lambda_\sigma^{\min}$	$\lambda_\sigma^{\max}$	$\lambda_{\min}^*$
5	113.19	236.49	113.11	113.26	235.89
6	260.82	362.10	260.72	260.86	361.34
7	449.55	524.70	449.45	449.60	523.16
8	679.45	728.36	679.04	679.55	724.50
9	950.28	976.66	949.58	950.49	969.81
10	1261.79	1272.09	1260.71	1262.23	1268.48
11	1613.78	1615.77	1610.89	1615.30	
12	2006.09	2006.11	1997.53	2010.41	
13	2438.60	2438.60	2403.42	2457.15	
14	2911.21	2911.21	2843.32	2947.17	
15	3423.83	3423.83	3260.54	3514.51	
16	3976.40	3976.40	3597.37	4211.88	

$v_1 = 4$ . The remaining part of the proof of Theorem 7 is divided into two parts. First, in section 6 a rigorous bound on the location of the unstable manifold close to the stationary point is obtained by analytical methods. Then the intersection of the manifold with the hyperplane and its location are proved by a computer assisted algorithm; see section 9. The following definition explains exactly what we mean by a computer assisted proof.

DEFINITION 8. *A proof is called computer assisted if it consists in finitely many elementary operations, but their number is so large that, although each step may be written down explicitly, it is only practical to perform such operations with a computer.*

We believe that a weakly singular solution exists in any dimension  $n \geq 5$ , but since our type of proof requires a finite number of steps for each value of  $n$ , we cannot prove this conjecture. We performed the computer assisted proof for  $n = 5, \dots, 16$  because the “interesting” phenomena of  $(P_\lambda)$  arise in these dimensions.

We expect the “singular parameter”  $\lambda_\sigma$  and the singular solution to be unique. However, also for this statement, we do not yet have a proof. See Open Problem 3 in section 8 below.

Table 1 summarizes our results:  $\lambda^*$  and  $\lambda_\sigma$  are the best, purely numerical, estimates for the values, up to two decimal places, while the numbers  $\lambda_\sigma^{\min}$ ,  $\lambda_\sigma^{\max}$ , and  $\lambda_{\min}^*$  are rigorously computed values as stated in Theorem 7.

Remark 9. We point out that both the approximate numerical computation and the computation with rigorous estimate on the error for  $\lambda^*$  become very difficult as  $n$  increases. For this reason the best rigorous estimate we have on  $\lambda_{\min}^*$  for  $n \geq 11$  is nothing but for  $\lambda_\sigma^{\min}$ , while the best numerical estimate we have on  $\lambda^*$  for  $n \geq 13$  is  $\lambda_\sigma$ . These values of  $n$  may be improved with a more accurate algorithm, but we do not feel that this would lead to a qualitative improvement of the result.

From Table 1 we immediately get the following.

COROLLARY 10. *For all  $n = 5, \dots, 10$  we have  $\lambda_\sigma < \lambda^*$ .*

Remark 11. We have numerical evidence that  $\lambda_\sigma < \lambda^*$  for  $n = 11, 12$  as well, but  $\lambda^* - \lambda_\sigma$  is much smaller than the rigorous estimate of the numerical error, and thus we do not have a proof. For  $n \geq 13$  the values of  $\lambda_\sigma$  and  $\lambda^*$  are closer than the numerical error; therefore we cannot even provide a conjecture supported by numerical evidence. If one could show uniqueness of the singular parameter  $\lambda_\sigma$  and that in fact  $\lambda_\sigma < \lambda^*$  in dimensions  $n \leq 12$ , one could conclude that here the extremal solution  $U_*$  is either regular or “strongly singular” (i.e.,  $\lim_{r \rightarrow 0} ru'(r)$  does not exist). For  $n \geq 13$  we expect the extremal solution  $U_*$  to be weakly singular. See Open Problems 3, 4, and 5 in section 8.




FIG. 1.

To complete the numerical inspection of the problem, we provide Figure 1, which shows the (regular) solution for  $n = 5$  and  $\lambda = \lambda^*$ .

Theorem 6 states that any weakly singular solution  $u = u(r)$  of (4) corresponds to a (weakly singular) solution  $v = v(s)$  of (6) which satisfies  $v(s) \approx -4s$  as  $s \rightarrow -\infty$ : this is because  $\bar{v}(s) = -4s$  is precisely the stationary point  $P_2$ . Hence, as a further consequence of Theorem 6, we have that any weakly singular solution  $U_\sigma$  behaves asymptotically like  $-4 \log r$  as  $r \rightarrow 0$ . Moreover, as may be checked by a simple calculation, the function  $r \mapsto -4 \log r$  solves the equation and the first boundary condition in  $(P_\lambda)$  for  $\lambda = 8(n - 2)(n - 4)$  but *not* the second boundary condition (recall also  $\lambda_\sigma > 8(n - 2)(n - 4)$  by Theorem 7). Contrary to what happens for the second order equation (1), the explicit form of the radial weakly singular solution does not seem simple to be determined; see also Proposition 34 below. To this end, we characterize it further by means of the following theorem.

**THEOREM 12.** *Let  $U_\sigma$  be a weakly singular solution with  $\lambda_\sigma > 8(n - 2)(n - 4)$  as it is obtained in Theorem 7 for  $5 \leq n \leq 16$ . Then*

$$U_\sigma(r) = -4 \log r + W(r),$$

where  $W$  is a bounded function satisfying

$$\lim_{r \rightarrow 0} W(r) = W_0 := \log \frac{8(n - 2)(n - 4)}{\lambda_\sigma} < 0$$

and (at least) one of the two following facts holds true:

- (i)  $W(r) - W_0$  changes sign infinitely many times in any neighborhood of  $r = 0$ .
- (ii)  $W(r) \geq \max[W_0, 2r^2 - 2]$  for all  $r \in (0, 1]$ .

If  $n \geq 13$ , case (ii) necessarily occurs.

Finally, the function  $W(r)$  is not analytic, i.e., not a convergent power series in  $r^2$  close to  $r_0 = 0$ .

*Remark 13.* It is quite surprising that the asymptotic behavior of weakly singular solutions of  $(P_\lambda)$  is the same as that of the quasi-linear equation  $-\Delta_4 u = \lambda e^u$ ; see [GPP]. Here  $-\Delta_p$  denotes the  $p$ -Laplace operator.

We conclude this section with a short remark concerning the behavior of  $(P_\lambda)$  in low dimensions.

*Remark 14.* In dimensions  $1 \leq n \leq 4$  the problem is subcritical and has a different behavior. In particular, there are no singular solutions.

The minimal solution is constructed as in the present paper. There is a parameter  $\lambda^* > 0$  such that for any  $\lambda \in (0, \lambda^*)$  there is precisely one minimal stable positive solution. Taking this one as a “trivial” solution, with the help of variational techniques (which apply only in a subcritical setting) one finds a second positive “large” solution above the minimal solution and unstable. For  $\lambda > \lambda^*$  there is no positive solution. Concerning the bifurcation diagram, one expects a smooth branch emanating from 0, extending until  $\lambda^*$ , where it bends back and approaches  $\lambda = 0$ , while the  $L^\infty$  norm of the solutions blows up. See also [We, Wi].

**3. Characterization of regular and weakly singular radial solutions.** In this section we perform a phase space analysis for the system (8), which corresponds to the radial version of  $(P_\lambda)$ . This gives some insight into which behavior of regular and of weakly singular radial solutions may be expected in dependence on the space dimension. These results are essential for the proofs of Theorems 7 and 12. For the proofs of Theorems 3 and 4 one may skip directly to sections 4 and 5.

**3.1. Analysis of the stationary points.** It is easy to verify that system (8) has only two stationary points:

$$P_1 = (4, 0, 0, 0) \quad \text{and} \quad P_2 = (0, 4n - 8, 16 - 8n, -8(n - 2)(n - 4)).$$

In order to linearize (8) in a neighborhood of  $P_1$ , we must just replace  $(8)_4$  with

$$v'_4(s) = 4v_4(s).$$

Then the linearized system has two distinct positive eigenvalues,  $\mu_1 = 2, \mu_2 = 4$ , and two distinct negative ones,  $\mu_3 = 2 - n, \mu_4 = 4 - n$ . We conclude that  $P_1$  is a hyperbolic point independently of the dimension.

Eigenvectors corresponding to the positive eigenvalues  $\mu_1, \mu_2$  in the neighborhood of  $P_1$  have the form

$$\alpha_1(1, -n, 0, 0) \quad \text{and} \quad \alpha_2(-1, n + 2, 2n + 4, 2n^2 + 4n),$$

where  $\alpha_1, \alpha_2 \in \mathbb{R} \setminus \{0\}$ . Therefore, the tangent hyperplane to the unstable manifold of  $P_1$  consists of those points in  $\mathbb{R}^4$  whose coordinates can be represented as

$$(10) \quad (\alpha_1 - \alpha_2, -n\alpha_1 + (n + 2)\alpha_2, (2n + 4)\alpha_2, (2n^2 + 4n)\alpha_2)$$

with  $\alpha_1, \alpha_2 \in \mathbb{R}$ .

Similarly, the tangent hyperplane to the stable manifold of  $P_1$  is spanned by eigenvectors corresponding to negative eigenvalues of the linearized system, that is,

$$\alpha_3(1, 0, 0, 0) \quad \text{and} \quad \alpha_4(1, -2, 2n - 4, 0),$$

where  $\alpha_3, \alpha_4 \in \mathbb{R} \setminus \{0\}$ .

Now consider the second critical point  $P_2$  of (8). In its neighborhood the linear approximation of  $(8)_4$  (the only nonlinear equation) takes the form

$$v'_4(s) = -8(n - 2)(n - 4)v_1(s).$$

Therefore, the eigenvalues of the linearized system in the neighborhood of  $P_2$  are the solutions of the fourth order algebraic equation

$$\nu(\nu - 2)(\nu + n - 2)(\nu + n - 4) - 8(n - 2)(n - 4) = 0;$$

hence,

$$\nu_{1,2,3,4} = \frac{1}{2} \left( 4 - n \pm \sqrt{M_1(n) \pm M_2(n)} \right),$$

where  $M_1(n) = n^2 - 4n + 8 = (n - 2)^2 + 4 > (n - 2)^2$  and  $M_2(n) = 4\sqrt{68 - 52n + 9n^2}$ . Therefore,

$$\nu_1 = \frac{1}{2} \left( 4 - n + \sqrt{M_1(n) + M_2(n)} \right) \quad \text{and} \quad \nu_2 = \frac{1}{2} \left( 4 - n - \sqrt{M_1(n) + M_2(n)} \right)$$

are real numbers. It is easy to see that

$$\nu_2 < 0 < \nu_1 \quad \text{for all } n \geq 4.$$

Moreover, for  $5 \leq n \leq 12$ , we have  $M_1(n) - M_2(n) < 0$ , while for  $n \geq 13$  there holds  $M_1(n) - M_2(n) > 0$ . Therefore, for  $5 \leq n \leq 12$  the eigenvalues

$$\nu_3 = \frac{1}{2} \left( 4 - n + \sqrt{M_1(n) - M_2(n)} \right) \quad \text{and} \quad \nu_4 = \frac{1}{2} \left( 4 - n - \sqrt{M_1(n) - M_2(n)} \right)$$

are complex conjugate with the real part

$$\operatorname{Re} \nu_3 = \operatorname{Re} \nu_4 = \frac{1}{2}(4 - n) < 0,$$

while for  $n \geq 13$  both  $\nu_3$  and  $\nu_4$  are real,  $\nu_3 < 0$  and  $\nu_4 < 0$ .

This analysis implies that for all  $n \geq 5$  the critical point  $P_2$  of system (8) is also hyperbolic, but its stable manifold is three-dimensional and the unstable manifold is one-dimensional. Moreover, taking into account that for  $5 \leq n \leq 12$  we have  $\operatorname{Im}\nu_3 = -\operatorname{Im}\nu_4 \neq 0$ , we deduce from the general theory of critical points (see, for example, [A2]) that for these values of  $n$  (and only for them) trajectories in the stable manifold of  $P_2$  locally have the form of a spiral.

**3.2. Proof of Theorem 6.** We first consider regular solutions. It will prove to be useful to have the following meaning of  $v_1, \dots, v_4$  in terms of derivatives of  $u$  in mind:

$$(11) \quad \begin{cases} v_1(s) = e^s u'(e^s) + 4, \\ v_2(s) = -e^{2s} \cdot \Delta u(e^s), \\ v_3(s) = -e^{3s} (\Delta u)'(e^s), \\ v_4(s) = -\lambda e^{4s} e^{u(e^s)}. \end{cases}$$

If  $u$  is a regular solution of  $(P_\lambda)$ , then  $u, u', \Delta u$ , and  $(\Delta u)'$  stay bounded in particular for  $r \searrow 0$ , i.e., for  $s \rightarrow -\infty$ . So, we get immediately from (11) the first part of the statement.

To prove the converse, assume that

$$\lim_{s \rightarrow -\infty} (v_1(s), v_2(s), v_3(s), v_4(s)) = P_1$$

so that

$$(12) \quad \lim_{r \searrow 0} ru'(r) = \lim_{r \searrow 0} r^2 \Delta u(r) = \lim_{r \searrow 0} r^3 (\Delta u)'(r) = \lim_{r \searrow 0} r^4 e^{u(r)} = 0.$$

The first limit yields particularly, that for  $r > 0$  small enough,

$$u(r) \leq -\frac{1}{2} \log(r), \quad e^{u(r)} \leq r^{-1/2}.$$

Using the differential equation  $(P_\lambda)$  and the growth conditions (12) (observe  $n > 4$ ), we obtain successively for  $r$  close to 0

$$(\Delta u)'(r) = O(r^{1/2}), \quad \Delta u(r) = O(1), \quad u'(r) = O(r), \quad u(r) = O(1).$$

That means that  $u$  is regular.

Next, we characterize weakly singular solutions. All the limits are intended as  $s \rightarrow -\infty$ ; with  $c$  we denote generic constants.

Note first that if  $\lim V(s) = P_2$ , then the solution is weakly singular.

In order to prove the converse, we claim that

$$(13) \quad v'(s) \rightarrow -4.$$

To this end, we exclude all the other cases; recall that  $\lim v'(s)$  exists by definition of weakly critical solutions.

(A) It cannot be that  $\lim v'(s) = c \in (-\infty, -4)$ .

For contradiction, if  $\lim v'(s) = c < -4$ , then by  $(7)_1$  we infer

$$(14) \quad \lim v_1(s) = c + 4 < 0,$$

and by  $(7)_4$  we get

$$(15) \quad v_4(s) \rightarrow -\infty.$$

Write  $(8)_3$  as

$$\frac{d}{ds}[e^{(n-4)s}v_3(s)] = e^{(n-4)s}v_4(s)$$

so that by (15) we infer that the map  $s \mapsto e^{(n-4)s}v_3(s)$  is decreasing in a neighborhood of  $-\infty$ , and therefore it admits a limit. If  $e^{(n-4)s}v_3(s) \rightarrow c \geq 0$ , then by  $(8)_3$  and (15) we get  $v'_3(s) \rightarrow -\infty$  and hence  $v_3(s) \rightarrow +\infty$ . If  $e^{(n-4)s}v_3(s) \rightarrow c < 0$ , then  $v_3(s) \rightarrow -\infty$ . In any case we obtain

$$(16) \quad |v_3(s)| \rightarrow +\infty.$$

A completely similar (but slightly more involved) argument shows that  $(8)_2$  and (16) entail

$$(17) \quad |v_2(s)| \rightarrow +\infty.$$

Finally,  $(8)_1$ , (14), and (17) furnish  $|v'_1(s)| \rightarrow +\infty$ , which contradicts (14).

(B) It cannot be that  $v'(s) \rightarrow -\infty$ .

For contradiction, assume that  $v'(s) \rightarrow -\infty$ : then by  $(7)_1$  we have

$$(18) \quad v_1(s) \rightarrow -\infty,$$

and by  $(7)_4$  we get

$$(19) \quad v_4(s) \rightarrow -\infty;$$

moreover,

$$(20) \quad \frac{v(s)}{s} \rightarrow -\infty.$$

We may rewrite (8)<sub>3</sub> as

$$\frac{d}{ds}[e^{(n-4)s}v_3(s)] = e^{(n-4)s}v_4(s) = -\lambda e^{ns+v(s)} \rightarrow -\infty,$$

where the second equality is just (7)<sub>4</sub> and the infinite limit is a consequence of (20): the previous limit yields  $e^{(n-4)s}v_3(s) \rightarrow +\infty$  and, in turn,

$$(21) \quad v_3(s) \rightarrow +\infty.$$

Similarly, we may rewrite (8)<sub>2</sub> as

$$\frac{d}{ds}[e^{-2s}v_2(s)] = e^{-2s}v_3(s) \rightarrow +\infty,$$

where the infinite limit is a consequence of (21): hence, we deduce that  $e^{-2s}v_2(s) \rightarrow -\infty$ , which, together with (8)<sub>2</sub> and (21), shows that  $v_2(s) \rightarrow -\infty$ . Inserting this into (7)<sub>2</sub> gives  $v''(s) + (n-2)v'(s) \rightarrow +\infty$ , and therefore  $v'(s) + (n-2)v(s) \rightarrow -\infty$ : hence,

$$\text{there exists } \sigma < 0 \text{ such that } v'(s) + (n-2)v(s) < 0 \quad \text{for all } s \leq \sigma.$$

We rewrite this inequality as

$$\frac{d}{ds}[e^{(n-2)s}v(s)] < 0 \quad \text{for all } s \leq \sigma;$$

integrating it over  $[s, \sigma]$  and taking into account that  $v(\sigma) > 0$ , we infer that

$$\text{there exists } K > 0 \text{ such that } v(s) \geq Ke^{(2-n)s} \quad \text{for all } s \leq \sigma.$$

Using (5) and returning to the function  $u$  (solution of  $(P_\lambda)$  and (4)), this shows that

$$\text{there exists } K > 0 \text{ such that } u(r) \geq \frac{K}{r^{n-2}} \quad \text{for all } r \leq e^\sigma;$$

this contradicts  $e^u \in L^1(B)$ .

(C) It cannot be that  $\lim v'(s) = c \in (-4, 0]$ .

For contradiction, if  $\lim v'(s) = c \in (-4, 0]$ , then by (7)<sub>1</sub> we infer

$$(22) \quad \lim v_1(s) = c + 4 > 0,$$

and by (7)<sub>4</sub> we get

$$(23) \quad v_4(s) \rightarrow 0.$$

Then from (8)<sub>3</sub> we deduce

$$(24) \quad v_3(s) \rightarrow 0,$$

because otherwise we would get a contradiction similar to that of case (A). Next, from (8)<sub>2</sub> and (24) we obtain

$$(25) \quad v_2(s) \rightarrow 0.$$

Since, by assumption,  $v_1$  has a limit, we deduce that necessarily  $v_1(s) \rightarrow 4$ . This, together with (23), (24), and (25), contradicts part (i) proved above.

By (A), (B), (C), statement (13) is proved. This shows that  $v_1(s) \rightarrow 0$ : inserting this into  $(8)_1$  gives  $v_2(s) \rightarrow 4(n - 2)$ . Inserting the latter into  $(8)_2$  yields  $v_3(s) \rightarrow -8(n - 2)$ ; finally, inserting this into  $(8)_3$  gives  $v_4(s) \rightarrow -8(n - 2)(n - 4)$ . This completes the proof of (ii).  $\square$

*Remark 15.* If in case (ii) of Theorem 6 we do not assume that  $\lim v'(s)$  exists, then we can merely show that  $\liminf v'(s) \leq -4 \leq \limsup v'(s)$ . Clearly, if one could prove that both inequalities are in fact equalities, then we would again have (13).

**4. Proof of Theorem 3.** We denote by  $\mathcal{K}$  the cone of nonnegative  $L^2$ -functions in  $B$ ,

$$\mathcal{K} = \{u \in L^2(B); u(x) \geq 0 \text{ for almost every } x \in B\},$$

and (for the sake of completeness) we prove the following weak formulation of Boggio's positivity preserving property [B], which we extensively use.

LEMMA 16. *Assume that  $u \in L^2(B)$  satisfies*

$$\int_B u \Delta^2 v \geq 0 \quad \text{for all } v \in \mathcal{K} \cap H^4 \cap H_0^2(B);$$

then  $u \in \mathcal{K}$ . Moreover, one has either  $u \equiv 0$  or  $u > 0$  a.e. in  $B$ .

*Proof.* (i) Take any  $\varphi \in \mathcal{K} \cap C_c^\infty(B)$  and let  $v_\varphi$  be the unique (classical) solution of

$$\begin{cases} \Delta^2 v_\varphi = \varphi & \text{in } B, \\ v_\varphi = \frac{\partial v_\varphi}{\partial \mathbf{n}} = 0 & \text{on } \partial B. \end{cases}$$

Then, by the classical Boggio principle [B], we infer that  $v_\varphi \in \mathcal{K}$ . Hence,  $v_\varphi$  is a possible test function for all  $\varphi$  so chosen, and therefore

$$\int_B u \varphi = \int_B u \Delta^2 v_\varphi \geq 0 \quad \text{for all } \varphi \in \mathcal{K} \cap C_c^\infty(B).$$

This shows that  $u \in \mathcal{K}$ .

(ii) By (i) we know that  $u \in \mathcal{K}$ . So, assume that  $u \not\equiv 0$  a.e. in  $B$  and let  $\phi$  denote the characteristic function of the set  $\{x \in B; u(x) = 0\}$  so that  $\phi \geq 0$ ,  $\phi \not\equiv 0$ . Let  $v_0$  be the unique (a.e.) solution of the problem

$$\begin{cases} \Delta^2 v_0 = \phi & \text{in } B, \\ v_0 = \frac{\partial v_0}{\partial \mathbf{n}} = 0 & \text{on } \partial B. \end{cases}$$

Then

$$v_0 \in \left( \bigcap_{q \geq 1} W^{4,q}(B) \right) \subset C^3(\bar{B})$$

and by Boggio's principle [B] we have  $v_0 > 0$  in  $B$ . By the biharmonic analogue of Hopf's lemma in balls (see [GS, Theorem 3.2], which also holds if  $\Delta^2 v_0 \in L^p(B)$  for some  $p > n/2$ ), we necessarily have  $\Delta v_0 > 0$  on  $\partial B$ . This last inequality allows us to

state that for all  $v \in C^4(\overline{B}) \cap H_0^2(B)$  there exists  $t_1 \leq 0 \leq t_0$  such that  $v + t_0v_0 \geq 0$  and  $v + t_1v_0 \leq 0$  in  $B$ . This, combined with the fact that

$$\int_B u \Delta^2 v_0 = \int_{\{u=0\}} u = 0,$$

enables us to show that both

$$0 \leq \int_B u \Delta^2(v + t_0v_0) = \int_B u \Delta^2 v \quad \text{and} \quad 0 \geq \int_B u \Delta^2(v + t_1v_0) = \int_B u \Delta^2 v.$$

Hence, we have for all  $v \in C^4(\overline{B}) \cap H_0^2(B)$

$$\int_B u \Delta^2 v = 0.$$

We need to show that  $C^4(\overline{B}) \cap H_0^2(B)$  is dense in  $H^4 \cap H_0^2(B)$ . For this purpose, take any function  $U \in H^4(B) \cap H_0^2(B)$  and put  $f := \Delta^2 U$ . We approximate  $f$  in  $L^2(B)$  by  $C^\infty(\overline{B})$ -functions  $f_k$  and solve  $\Delta^2 U_k = f_k$  in  $B$  under homogeneous Dirichlet boundary conditions. We then even have  $U_k \in C^\infty(\overline{B})$ , and by  $L^2$ -theory there holds  $\|U_k - U\|_{H^4(B)} \rightarrow 0$  as  $k \rightarrow \infty$ .

By the previous statement we may now conclude that

$$\text{for all } v \in H^4 \cap H_0^2(B) : \quad \int_B u \Delta^2 v = 0.$$

Since  $u \in L^2(\Omega)$ , we may take as  $v \in H^4 \cap H_0^2(B)$  the solution of  $\Delta^2 v = u$  under homogeneous Dirichlet boundary conditions. This finally yields  $u \equiv 0$ .  $\square$

In particular, thanks to Lemma 16 we may establish a result in the spirit of [BCMR], as follows.

LEMMA 17. *For all  $f \in L^1(B)$  such that  $f \geq 0$  a.e. in  $B$  there exists a unique  $u \in L^1(B)$  such that  $u \geq 0$  a.e. in  $B$  and which satisfies*

$$\int_B u \Delta^2 v = \int_B f v \quad \text{for all } v \in C^4(\overline{B}) \cap H_0^2(B);$$

moreover, there exists  $C > 0$  (independent of  $f$ ) such that  $\|u\|_1 \leq C \|f\|_1$ .

*Proof.* Uniqueness follows by means of the observation that  $L^\infty$ -functions may be approximated by a pointwise convergent but uniformly bounded sequence of  $C_c^\infty(B)$ -functions. This is applied to truncations of  $u$ , and suitable test functions  $v$  are obtained from approximations of the truncations of  $u$  by solving the biharmonic Dirichlet problems.

Existence follows by truncating  $f$  and by arguing as in the proof of [BCMR, Lemma 1], the only difference being the positivity preserving property, which is standard for the Laplacian; in our case we invoke Lemma 16.  $\square$

Combining the method of proof of Lemmas 16 and 17, one also has the following.

LEMMA 18. *Assume that  $u \in L^1(B)$  satisfies*

$$\int_B u \Delta^2 v \, dx \geq 0 \quad \text{for all } v \in \mathcal{K} \cap C^4(\overline{B}) \cap H_0^2(B);$$

then  $u \geq 0$  a.e. in  $B$ .

As was pointed out to us by Anna Dall’Acqua (TU Delft), similar techniques and the application of Weierstraß’s approximation theorem yield that also for the stronger conclusion of Lemma 16 it is enough to require  $u \in L^1(B)$ .

The previous lemmas enable us to make use of the super-solutions method, as follows.

LEMMA 19. *Let  $\lambda > 0$  and assume that there exists  $\bar{u} \in \mathcal{K}$  such that  $e^{\bar{u}} \in L^1(B)$  and*

$$\int_B \bar{u} \Delta^2 v \geq \lambda \int_B e^{\bar{u}} v \quad \text{for all } v \in \mathcal{K} \cap W^{4,p} \cap H_0^2(B).$$

*Then there exists a solution  $u$  of  $(P_\lambda)$  such that  $0 \leq u \leq \bar{u}$  a.e. in  $B$ .*

*Proof.* Let  $u_0 = \bar{u}$ , and for all  $m \in \mathbb{N}$ , define inductively the function  $u_{m+1}$  as the unique solution of

$$(26) \quad \int_B u_{m+1} \Delta^2 v = \lambda \int_B e^{u_m} v \quad \text{for all } v \in W^{4,p} \cap H_0^2(B).$$

Note that by Lemmas 16 to 18 the sequence  $\{u_m\}$  is well-defined and

$$u_m \in \mathcal{K}, \quad e^{u_m} \in L^1(B), \quad 0 \leq u_{m+1}(x) \leq u_m(x) \text{ for almost every } x \in B$$

for all  $m \in \mathbb{N}$ .

Since this sequence is pointwise decreasing, there exists  $u \in \mathcal{K}$  such that  $e^u \in L^1(B)$  and which is the pointwise limit of  $\{u_m\}$ . Then, letting  $m \rightarrow \infty$  in (26) and applying Lebesgue’s theorem, we obtain the result.  $\square$

Define  $\Lambda := \{\lambda \geq 0; (P_\lambda) \text{ admits a solution}\}$  and

$$\lambda^* := \sup \Lambda;$$

clearly  $0 \in \Lambda$  and so  $\Lambda \neq \emptyset$ . Moreover, by the implicit function theorem we know that  $\lambda^* > 0$ . It follows directly from Lemma 19 that  $\Lambda$  is an interval.

Let  $\lambda \in \Lambda$ ; then there exists  $u_\lambda$  satisfying (2). Taking into account that  $e^s \geq es$  for all  $s \geq 0$  with strict inequality whenever  $s \neq 1$ , and choosing  $v = \phi_1$  (the normalized positive first eigenfunction of (3)) as a test function in (2), we get

$$\lambda_1 \int_B u_\lambda \phi_1 = \int_B u_\lambda \Delta^2 \phi_1 = \lambda \int_B e^{u_\lambda} \phi_1 > \lambda e \int_B u_\lambda \phi_1,$$

which proves that

$$(27) \quad \lambda < \frac{\lambda_1}{e} \quad \text{for all } \lambda \in \Lambda.$$

We now prove the most delicate part of Theorem 3, namely, that for any  $\lambda < \lambda^*$ , there exists a *regular* solution.

LEMMA 20. *Assume that for some  $\mu > 0$  there exists a (possibly singular) solution  $u_0$  of  $(P_\mu)$ . Then for all  $0 < \lambda < \mu$  there exists a regular solution of  $(P_\lambda)$ .*

*Proof.* Let  $0 < \lambda < \mu$  and consider the (unique) functions  $u_1, u_2 \in L^1(B)$  satisfying, respectively,

$$\int_B u_1 \Delta^2 v = \lambda \int_B e^{u_0} v \quad \text{for all } v \in W^{4,p} \cap H_0^2(B),$$

$$(28) \quad \int_B u_2 \Delta^2 v = \lambda \int_B e^{u_1} v \quad \text{for all } v \in W^{4,p} \cap H_0^2(B).$$

Such functions exist by Lemma 17 and also belong to  $L^2(B)$ , since by Lemma 18 we have

$$(29) \quad u_0 > \frac{\lambda}{\mu} u_0 = u_1 \geq u_2 \quad \text{almost everywhere in } B.$$

Let  $\varphi(x) = (1 - |x|^2)^2$ ; it is readily verified that

$$(30) \quad \varphi \in H_0^2(B), \quad \Delta^2 \varphi = 8n(n+2).$$

We also need the following elementary statement:

$$(31) \quad \begin{aligned} &\text{for all } \vartheta > 1 \text{ and } \delta > 0 \text{ there exists } \gamma > 0 \\ &\text{such that } e^{\vartheta s} + \gamma - (1 + \delta)e^s \geq 0 \text{ for all } s \geq 0. \end{aligned}$$

Take  $\vartheta = \mu/\lambda$ ,  $\delta = n\lambda/4\mu$  and choose  $k > 0$  in such a way that

$$(32) \quad e^{\frac{\mu}{\lambda}s} + \frac{8n(n+2)}{\lambda} k \geq (1 + \delta)e^s \quad \text{for all } s \geq 0;$$

this choice is clearly allowed by (31). Thanks to (30) and (32) we find

$$\begin{aligned} \int_B (u_1 + k\varphi)\Delta^2 v &= \int_B [\lambda e^{u_0} + 8n(n+2)k]v = \int_B [\lambda e^{\frac{\mu}{\lambda}u_1} + 8n(n+2)k]v \\ &\geq \lambda(1 + \delta) \int_B e^{u_1} v = (1 + \delta) \int_B u_2 \Delta^2 v \\ &\quad \text{for all } v \in \mathcal{K} \cap W^{4,p} \cap H_0^2(B). \end{aligned}$$

Hence, by Lemma 16 we infer that  $u_2 \leq \frac{u_1+k\varphi}{1+\delta}$  in  $B$ ; in particular, we get

$$e^{u_2} \leq e^{\frac{k}{1+\delta}\varphi} e^{\frac{\lambda}{\mu(1+\delta)}u_0},$$

from which we get at once that

$$(33) \quad e^{u_2} \in L^{\frac{n}{4} + \frac{\mu}{\lambda}}(B)$$

since  $\varphi \in L^\infty(B)$  and  $e^{u_0} \in L^1(B)$  (recall also our choice of  $\delta$ ). Finally, consider  $u_3 \in L^2(B)$  such that

$$\int_B u_3 \Delta^2 v = \lambda \int_B e^{u_2} v \quad \text{for all } v \in W^{4,p} \cap H_0^2(B).$$

By (33) and elliptic regularity [ADN], we deduce that

$$u_3 \in W^{4, \frac{n}{4} + \frac{\mu}{\lambda}}(B) \subset L^\infty(B).$$

Moreover, by (28), (29), and Lemma 16 we infer that  $u_3 \leq u_2$  and hence

$$\int_B u_3 \Delta^2 v \geq \lambda \int_B e^{u_3} v \quad \text{for all } v \in \mathcal{K} \cap W^{4,p} \cap H_0^2(B).$$

We have so found a weak bounded supersolution  $u_3$  of  $(P_\lambda)$ , and the statement follows from Lemma 19.  $\square$

With the help of Lemma 20 we can now show the following.

LEMMA 21. *For all  $0 \leq \lambda < \lambda^*$ , the minimal solution  $U_\lambda$  exists, is regular, and is radially symmetric.*

*Proof.* By the preceding lemma we have the existence of a regular solution  $u_\lambda$  of  $(P_\lambda)$ . This may serve as a (classical) supersolution of  $(P_\lambda)$ , while  $U_0 \equiv 0$  is a subsolution. Hence, the minimal solution  $U_\lambda$  of  $(P_\lambda)$  may be obtained as the increasing limit of the following sequence  $\{U_m\}$ :

$$\begin{cases} \Delta^2 U_{m+1} = \lambda e^{U_m} & \text{in } B, \\ U_{m+1} = \frac{\partial U_{m+1}}{\partial \mathbf{n}} = 0 & \text{on } \partial B \end{cases} \quad (m \geq 0).$$

Since  $U_0$  is radially symmetric, so is  $U_1$ ; similarly, all the functions  $U_m$  are radially symmetric: therefore, their (pointwise) limit  $U_\lambda$  is also radially symmetric.  $\square$

The previous lemma allows us to show that the interval  $\Lambda$  is closed: we first remark that the map  $\lambda \mapsto U_\lambda(x)$  is strictly increasing for all  $x \in B$  (in view of Lemma 16). If  $0 \leq \lambda < \mu < \lambda^*$ , the minimal solution  $U_\mu$  of  $(P_\mu)$  is a (strict) supersolution for  $(P_\lambda)$ . Therefore

$$(34) \quad U_*(x) := \lim_{\lambda \rightarrow \lambda^*} U_\lambda(x) \in [0, \infty]$$

exists for all  $x \in B$ . In fact, more can be said about this limit, as follows.

LEMMA 22. *Let  $U_*$  be the function defined in (34). Then  $U_*(x)$  is finite for almost every  $x \in B$  and  $U_*$  solves  $(P_\lambda)$  for  $\lambda = \lambda^*$ . Moreover,  $U_\lambda \rightarrow U_*$  in  $H_0^2(B)$  as  $\lambda \uparrow \lambda^*$ . Finally,  $U_*$  is radially symmetric.*

*Proof.* By Lemma 21 we have  $U_\lambda \in C^\infty(\bar{B})$ , and therefore, by using the generalized Pohozaev identity [P] by Pucci and Serrin [PS] and by arguing as in the proof of [GMP, Théorème 2], we obtain that the set  $\{U_\lambda; \lambda < \lambda^*\}$  is bounded in  $H_0^2(B)$ , and hence  $U_\lambda \rightharpoonup U_*$  in  $H_0^2(B)$ , up to a subsequence (this follows by uniqueness of the pointwise limit). This shows that  $U_*$  is a.e. finite, that  $U_*$  solves  $(P_\lambda)$  for  $\lambda = \lambda^*$ , and also that  $U_* e^{U_*} \in L^1(B)$ . Finally, since  $U_\lambda e^{U_\lambda} \leq U_* e^{U_*}$ , by Lebesgue's theorem we deduce that

$$\frac{1}{\lambda} \int_B |\Delta U_\lambda|^2 = \int_B U_\lambda e^{U_\lambda} \rightarrow \int_B U_* e^{U_*} = \frac{1}{\lambda^*} \int_B |\Delta U_*|^2 \quad \text{as } \lambda \uparrow \lambda^*,$$

which, together with weak convergence, shows that  $U_\lambda \rightarrow U_*$  in the norm topology of  $H_0^2(B)$ ; since the above arguments may be repeated for any sequence in  $\{U_\lambda; \lambda < \lambda^*\}$ , the result follows without extracting subsequences.

Finally, by Lemma 21, all the minimal solutions  $U_\lambda$  (for  $0 < \lambda < \lambda^*$ ) are radially symmetric. Then by (34) also  $U_*$  is radially symmetric.  $\square$

Remark 23. The proof of Lemma 22 may also be obtained by exploiting the stability of the minimal solution  $U_\lambda$  (see Proposition 37(i) below) and by arguing as in [BV, Remark 3.3].

Finally, we claim that

$$(35) \quad \lambda^* \geq 14.72(n-1)(n-3).$$

Indeed, this holds true by Lemma 19 since the function  $\bar{u}(x) = 7.36(1 - |x|)^2$  is a weak supersolution ( $\bar{u} \in C^\infty(\bar{B} \setminus \{0\})$ ) of  $(P_\lambda)$  for all  $\lambda \leq 14.72(n-1)(n-3)$ .

*Proof of Theorem 3.* The upper bound for  $\lambda^*$  follows from (27) and from Lemma 22, the latter saying that  $\lambda^* \in \Lambda$ . The lower bound for  $\lambda^*$  is proved in (35). Statement (i) follows from Lemmas 20 and 21. The map  $\lambda \mapsto U_\lambda(x)$  is nondecreasing for all  $x$  by Lemma 19 and strictly increasing by Lemma 16; this proves the first part of statement (ii). The second parts of (ii) and (iii) follow from Lemma 22. Finally, the radial symmetry of  $U_*$  and of all the minimal solutions  $U_\lambda$  (for  $\lambda < \lambda^*$ ) is obtained in Lemmas 22 and 21, respectively. The regular minimal solutions  $U_\lambda$  (for  $\lambda < \lambda^*$ ) are strictly radially decreasing in view of [So]. Passing to the limit, we also get that  $U^*$  is radially decreasing.  $\square$

*Remark 24.* The above analysis does not allow us to establish whether the extremal solution  $U_*$  is regular, weakly singular, or singular. However, since it is radially symmetric, in the regular and weakly singular case, Theorem 6 describes the behavior of  $U_*$  when studied in the phase space  $\mathbb{R}^4$ . With our computer assisted proof, we may then find some space dimensions where the first case certainly occurs, provided that we can also show uniqueness of the weakly singular solution and the corresponding parameter  $\lambda_\sigma$ .

**5. Proof of Theorem 4.** We first show that

$$(36) \quad U_\lambda \rightarrow 0 \quad \text{uniformly as } \lambda \rightarrow 0.$$

Since this is standard, we just briefly sketch its proof. By Theorem 3 we know that

$$0 < \lambda < \mu < \lambda^* \implies U_\lambda(x) < U_\mu(x) \quad \text{if } |x| < 1.$$

Then, by multiplying the equation in  $(P_\lambda)$  by  $U_\lambda$  and by integrating by parts, we obtain that  $\|U_\lambda\|_{H_0^2(B)}$  remains bounded. Hence, up to a subsequence,  $\{U_\lambda\}$  converges in the weak  $H_0^2(B)$  topology to  $U_0 \equiv 0$ , which is the unique solution of  $(P_0)$ . By convergence of the norms, we infer that the convergence is in the norm topology. Finally, by pointwise convergence and elliptic regularity, we infer (36).

Next, note that  $V_\lambda$  satisfies

$$(37) \quad \begin{cases} \Delta^2 V_\lambda = \lambda & \text{in } B, \\ V_\lambda = \frac{\partial V_\lambda}{\partial \mathbf{n}} = 0 & \text{on } \partial B. \end{cases}$$

Therefore,  $\Delta^2 U_\lambda > \Delta^2 V_\lambda$ , and the inequality  $U_\lambda > V_\lambda$  follows by Lemma 16.

In order to prove the last statement of Theorem 4, note that from (36) we infer

$$\text{for all } \varepsilon > 0 \quad \text{there exists } \lambda_\varepsilon > 0 \quad \text{such that} \quad \lambda < \lambda_\varepsilon \implies \|U_\lambda\|_\infty < \varepsilon.$$

So, fix  $\varepsilon > 0$  and let  $\lambda < \lambda_\varepsilon$ . Then (37) entails

$$\Delta^2 U_\lambda = \lambda e^{U_\lambda} < \lambda e^\varepsilon = e^\varepsilon \Delta^2 V_\lambda \quad \text{in } B.$$

This shows that  $U_\lambda(x) < e^\varepsilon V_\lambda(x)$  for all  $x \in B$ , and the result follows by arbitrariness of  $\varepsilon$ .

**6. Proof of Theorem 7.** The proof of Theorem 7 is obtained with computer assistance. We first describe the numerical procedure used to obtain the approximate values for  $\lambda_\sigma$  and  $\lambda^*$ ; then we show how the algorithm can be made rigorous. We maintain here the same notation as in section 3. The computation of  $\lambda_\sigma$  is somehow simpler than the computation of  $\lambda^*$ , since the unstable manifold of  $P_2$  is one-dimensional. We

choose a point  $\bar{v} = P_2 + r e_1$  where  $e_1$  is an eigenvector corresponding to the unstable manifold and  $r$  is some small value. We solve system (8) with  $\bar{v}$  as the initial condition and look for the intersection of the solution with the hyperplane  $v_1 = 4$ . The choice of a positive or negative  $r$  leads to a different result, since the manifold is made of two branches: it turns out that one branch never appears to intersect the hyperplane, while the other branch always does. If the solution intersects the hyperplane  $v_1 = 4$  at some point  $\hat{v} = (\hat{v}_1, \hat{v}_2, \hat{v}_3, \hat{v}_4)$  such that  $\hat{v}_4 < 0$ , by Theorem 6 and by (7) we have numerical evidence of a singular solution at  $\lambda = -\hat{v}_4$ .

In order to compute the value of  $\lambda^*$ , we have to study the two-dimensional unstable manifold of  $P_1$ . The direction on the tangent hyperplane can be parametrized by an angle  $\vartheta$ . In order to find the largest value for  $\lambda$ , we use a *directional shooting* method; i.e., we choose some value  $\vartheta$  (the shooting direction) and solve the equation with starting point  $\bar{v} = P_1 + r(e_1 \sin \vartheta + e_2 \cos \vartheta)$ , where  $e_1$  and  $e_2$  are the orthonormalized eigenvectors corresponding to the (tangent) unstable manifold and  $r > 0$  is some small arbitrarily chosen value. If the solution intersects the hyperplane  $v_1 = 4$  at some point  $\hat{v} = (\hat{v}_1, \hat{v}_2, \hat{v}_3, \hat{v}_4)$  such that  $\hat{v}_4 < 0$ , then by Theorem 6 and by (7) and (8) we have numerical evidence of a regular solution for  $\lambda = -\hat{v}_4$ . By varying  $\vartheta$  we can look for the maximal value of  $\lambda$ .

Of course these procedures do not lead to an exact value for two reasons. First, we can choose only  $\bar{v}$  on the unstable manifold of the linearized equation, and although we know that we are close to the manifold of the full equation, we are not exactly on it. Second, the algorithm used to solve the differential equation provides an accurate, but not rigorous, solution. We address the problem of proving that a branch of the unstable manifold of  $P_2$  does intersect the hyperplane  $v_1 = 4$  and of computing a rigorous estimate for the values  $\lambda_\sigma$  and  $\lambda^*$  in the following sections.

**6.1. Rigorous bounds for the manifolds.** We first address the general problem of computing rigorous bounds for the location of the unstable manifold in the neighborhood of a stationary hyperbolic point of an ODE. The same technique could be applied to the stable manifold as well, but in this paper we are not interested in it.

Let  $f \in C^2(\mathbb{R}^d, \mathbb{R}^d)$ ,  $d \geq 2$ . We consider the equation  $\dot{x} = f(x)$  and assume that 0 is a hyperbolic stationary point. Then

$$(38) \quad \dot{x} = Ax + N(x),$$

where

$$(39) \quad A = \nabla f(0), \quad N(x) = O(|x|^2) \text{ as } x \rightarrow 0$$

and all eigenvalues of  $A$  have nonzero real part. Let  $\varphi(x, t)$  be the flow induced by (38) and let  $\varphi_A(x, t)$  be the flow induced by the linear equation  $\dot{x} = Ax$ . Let  $S_0$  (resp.,  $U_0$ ) be the span of all eigenvectors corresponding to the eigenvalues with negative (resp., positive) real part.  $S_0$  (resp.,  $U_0$ ) is called the stable (resp., unstable) subspace, and it is characterized as follows:  $S_0$  (resp.,  $U_0$ ) is the set of points  $x \in \mathbb{R}^d$  such that  $\varphi_A(x, t) \rightarrow 0$  as  $t \rightarrow +\infty$  (resp.,  $t \rightarrow -\infty$ ). It is well known that the full equation also admits a stable manifold  $S$  (resp., an unstable manifold  $U$ ) still defined as the set of points  $x \in \mathbb{R}^d$  such that  $\varphi(x, t) \rightarrow 0$  as  $t \rightarrow +\infty$  (resp.,  $t \rightarrow -\infty$ ). Such a manifold is tangent at the origin to  $S_0$  (resp.,  $U_0$ ). If  $S_0$  (resp.,  $U_0$ ) is empty, then there exists a neighborhood of the origin which is a subset of  $U$  (resp.,  $S$ ). We are interested in the case when both manifolds are nontrivial, and we wish to study the intersection of the unstable manifold with some other manifold  $P$ . In order to achieve this goal,

we consider a point  $\bar{x} \in U \setminus \{0\}$  and study  $\varphi(\bar{x}, t)$ . If we can prove that  $\varphi(\bar{x}, t_0) \in P$  for some positive  $t_0$ , then we infer that  $U \cap P \neq \emptyset$ , and we also know the intersection point. The main problem to address is that the only point of the manifold we know precisely is the origin: the other points lie very close to  $U_0$ , at least in a neighborhood of 0, but we do not know their explicit position. We proceed as follows.

There exists an invertible matrix  $M$  such that  $B := M^{-1}AM$  is block diagonal, i.e., the canonical basis  $\{e_i\}$  of  $\mathbb{R}^d$  is split in  $S'_0 \cup U'_0$ , where  $S'_0 = \text{span}\{e_1, \dots, e_m\}$  is the stable eigenspace and  $U'_0 = \text{span}\{e_{m+1}, \dots, e_d\}$  is the unstable eigenspace. If we let  $y = M^{-1}x$ , the (38) can be written as

$$(40) \quad \dot{y} = By + M^{-1}N(My) =: g(y).$$

By (39), for all  $\varepsilon > 0$  there exists  $\beta > 0$  such that  $|N(x)| \leq \beta|x|^2$  for all  $|x| \leq \varepsilon$ . Let  $\alpha < 0$  be the maximum of the real parts of the eigenvalues with negative real parts,  $\gamma = -\frac{\alpha}{\beta m_1^2 m_2}$ ,  $m_1 = \|M\|$ , and  $m_2 = \|M^{-1}\|$ . Choose  $\varepsilon > 0$ ; let  $\beta > 0$  as above; choose a vector  $\hat{y} \in U'_0 \setminus \{0\}$  of norm  $r \leq \varepsilon$  and  $k > 1$ . Let  $P_s$  be the orthogonal projection onto  $S'_0$ ; let  $P_u$  be the orthogonal projection onto the linear space spanned by  $\hat{y}$ ; and let

$$(41) \quad \Xi = \left\{ y \in \mathbb{R}^d : \frac{\gamma}{k} |P_s y| \leq |P_u y|^2 \leq r^2 \right\}.$$

We show that, under a suitable choice of  $k > 1$  and  $0 < r \leq \varepsilon$ , for all  $y \in \partial\Xi$  such that  $|P_u y| < r$  the flow is inward; i.e., given  $\bar{y} \in \Xi$  we want the solution of the Cauchy problem  $\dot{y}(t) = g(y(t))$ ,  $y(0) = \bar{y}$  to leave  $\Xi$  only through the set  $\{y \in \partial\Xi : |P_u y| = r\}$ . If this happens, then for all  $\hat{y} \in U'_0$  satisfying  $|\hat{y}| = r$  either the unstable manifold intersects the set

$$(42) \quad \kappa := \kappa_{\hat{y}} := \hat{y} + \left\{ \tilde{y} \in S'_0 : |\tilde{y}| \leq \frac{k}{\gamma} r^2 \right\}$$

or it is entirely contained in  $\Xi$ . As a result, to study a branch of the unstable manifold it is sufficient to exclude the second case and consider the initial value problem for all  $\bar{y} \in \kappa$ .

LEMMA 25. *Choose  $\varepsilon > 0$  and  $k > 1$ . Let  $\alpha, \beta, \gamma, m_1, m_2$ , and  $\Xi$  be as above and let*

$$(43) \quad r = \min \left\{ \frac{\varepsilon\gamma}{m_1\sqrt{\gamma^2 + k^2}}, 1, \frac{\sqrt{k-1}}{k}\gamma, \frac{\gamma}{2} \right\}.$$

For all  $\bar{y} \in \partial\Xi$  such that  $0 < |P_u \bar{y}| < r$  we have

$$(44) \quad (g(\bar{y}), P_s \bar{y}) < 0.$$

*Proof.* Let  $\hat{y} = P_s \bar{y}$ ,  $\tilde{y} = P_u \bar{y}$ ,  $\hat{r} = |\hat{y}|$ , and  $\tilde{r} = |\tilde{y}|$ . Since  $\tilde{r} < r \leq \frac{\varepsilon\gamma}{m_1\sqrt{\gamma^2 + k^2}}$ , then  $|M\bar{y}| \leq \varepsilon$ , and therefore  $|N(\bar{y})| \leq \beta|\bar{y}|^2$ . We have

$$\begin{aligned} (B\bar{y}, \hat{y}) + (M^{-1}N(M\bar{y}), \hat{y}) &= (B\hat{y}, \hat{y}) + (N(M\bar{y}), (M^{-1})^t \hat{y}) \\ &\leq \alpha\hat{r}^2 + \beta m_1^2 m_2 (\hat{r}^2 + \tilde{r}^2)\hat{r}. \end{aligned}$$

Then a simple computation shows that (44) is implied by

$$(45) \quad \frac{1}{2} \left( \gamma + \sqrt{\gamma^2 - 4\tilde{r}^2} \right) > \hat{r} > \frac{1}{2} \left( \gamma - \sqrt{\gamma^2 - 4\tilde{r}^2} \right).$$

The first inequality is satisfied because  $\tilde{r} < r \leq \frac{\gamma}{2}$ . For all  $\bar{y} \in \partial\Xi$ ,  $0 < |P_u\bar{y}| < r$ , by (41), we have  $\hat{r} = \frac{k}{\gamma}\tilde{r}^2$  and

$$\frac{1}{2} \left( \gamma - \sqrt{\gamma^2 - 4\tilde{r}^2} \right) < k \frac{\tilde{r}^2}{\gamma} \quad \text{if } \tilde{r} < \frac{\sqrt{k-1}}{k} \gamma.$$

Since  $\tilde{r} < r$ , then (45) and therefore (44) hold.  $\square$

We need a condition which ensures that the invariant manifold is not entirely contained in  $\Xi$ , but it intersects  $\kappa$  at some point. Let  $\alpha'$  be the minimum of the real parts of the eigenvalues of  $B$  with positive real parts.

LEMMA 26. *If  $r, \alpha, \alpha', k$ , and  $\Xi$  are as above and*

$$(46) \quad \alpha' + \frac{\alpha}{\gamma} \left( r + \frac{k^2}{\gamma^2} r^3 \right) > 0,$$

*then there exists  $\delta > 0$  such that  $(g(\bar{y}), P_u\bar{y}) \geq \delta|P_u\bar{y}|^2$  for all  $\bar{y} \in \Xi$ , and therefore the component of the flow in the direction of the unstable manifold is always increasing in  $\Xi$ , together with its first derivative.*

*Proof.* Choose  $\delta > 0$  satisfying

$$\beta m_1^2 m_2 \left( r + \frac{k^2}{\gamma^2} r^3 \right) \leq \alpha' - \delta.$$

Fix  $\bar{y} \in \Xi$  and let  $\hat{y} = P_s\bar{y}$ ,  $\tilde{y} = P_u\bar{y}$ ,  $\hat{r} = |\hat{y}|$ , and  $\tilde{r} = |\tilde{y}|$ . By (46) and the definition of  $\gamma$ , such a  $\delta$  exists. We have

$$\begin{aligned} (B\tilde{y}, \tilde{y}) + (M^{-1}N(M\tilde{y}), \tilde{y}) &= (B\tilde{y}, \tilde{y}) + (N(M\tilde{y}), (M^{-1})^t\tilde{y}) \\ &\geq \alpha'\tilde{r}^2 - \beta m_1^2 m_2 (\hat{r}^2 + \tilde{r}^2) \tilde{r} \geq \delta\tilde{r}^2, \end{aligned}$$

because  $\hat{r} \leq \frac{k}{\gamma}\tilde{r}^2$  by the definition of  $\Xi$ .  $\square$

LEMMA 27. *Let  $r, \alpha, \alpha', k, \hat{y}$ , and  $\kappa$  be as above. The unstable manifold tangent to  $\hat{y}$  intersects  $\kappa$ .*

*Proof.* By Lemma 26 the unstable manifold cannot be entirely contained in  $\Xi$ . By Lemmas 25 and 26 it can only exit through  $\kappa$ .  $\square$

In the next subsection we apply these ideas in order to prove Theorem 7.

**6.2. The computer assisted proofs.** We apply the general result stated in the previous subsection to system (8).

We first consider the point  $P_1 = (4, 0, 0, 0)$ . Let  $x = v - P_1$ . System (8) takes the form (38) with

$$A_1 = \begin{bmatrix} 2-n & -1 & 0 & 0 \\ 0 & 2 & 1 & 0 \\ 0 & 0 & 4-n & 1 \\ 0 & 0 & 0 & 4 \end{bmatrix},$$

$N(x) = (0, 0, 0, x_1x_4)$ , and  $\alpha = 4 - n$ .

If we consider the linearization at  $P_2 = (0, 4n - 8, 16 - 8n, -8(n - 2)(n - 4))$  and set  $x = v - P_2$ , then system (8) can be written as (38) with

$$A_2 = \begin{bmatrix} 2-n & -1 & 0 & 0 \\ 0 & 2 & 1 & 0 \\ 0 & 0 & 4-n & 1 \\ -8(-4+n)(-2+n) & 0 & 0 & 0 \end{bmatrix},$$

and again  $N(x) = (0, 0, 0, x_1x_4)$ . From section 3.1 we know that if  $n = 5, \dots, 12$ , the eigenvalues are  $((4 - n)/2 + i\sigma, (4 - n)/2 - i\sigma, \lambda_1, \lambda_2)$ , where  $\lambda_1 < (4 - n)/2 < 0$  and  $\lambda_2 > 0$ . It turns out that  $\alpha = (4 - n)/2$ . If  $n \geq 13$ , all eigenvalues are real and

$$\alpha = 2 - \frac{1}{2}n + \frac{1}{2}\sqrt{8 - 4n + n^2 - 4\sqrt{68 - 52n + 9n^2}}.$$

We remark that since the nonlinear part is very simple, it is possible to obtain a better estimate for the coefficients  $\beta$ ,  $m_1$ , and  $m_2$  than the one we had in section 6.1.

In the following, let  $M$  be the matrix that diagonalizes either  $A_1$  or  $A_2$  and let  $|M_i|$  be the (Euclidean) norm of the  $i$ th row of  $M$ .

LEMMA 28. *For all  $y_1, y_2 \in \mathbb{R}^4$  the following inequality holds:*

$$(N(My_1), (M^{-1})^t y_2) \leq |M_1| |M_4| |(M^{-1})_4^t| |y_1|^2 |y_2|.$$

*Proof.* We have

$$(N(My_1), (M^{-1})^t y_2) = (My_1)_1 (My_1)_4 ((M^{-1})^t y_2)_4,$$

where we denoted by  $(Av)_i$  the  $i$ th component of the vector  $(Av)$ , i.e., the scalar product of the  $i$ th row of  $A$  with the vector  $v$ . The conclusion follows by the definition of  $|M_i|$ .  $\square$

By the above lemma we infer that  $\gamma$  may be obtained as

$$(47) \quad \gamma = -\frac{\alpha}{|M_1| |M_4| |(M^{-1})_4^t|}$$

and  $\varepsilon$  may be chosen arbitrarily.

To compute a rigorous enclosure  $[\lambda_\sigma^{\min}, \lambda_\sigma^{\max}]$  for the value of  $\lambda_\sigma$ , we fix  $n$  and compute the value  $\gamma$  in (47). We can choose  $k > 1$  and  $r > 0$  satisfying (43) and (46). We have some degree of arbitrariness: we prefer a small  $r$  in order to have a small set  $\kappa$ , but we also like a large  $r$  in order to reach the hyperplane in fewer time steps. It is also convenient to have the smallest possible  $k$ , since it also implies a smaller set  $\kappa$ . We have to make an empirical choice by trying different values and selecting the best trade-off. It turns out that it is convenient to choose  $r$  first, set

$$(48) \quad k = \frac{\gamma^2 - \sqrt{\gamma^4 - 4r^2\gamma^2}}{2r^2},$$

and check whether (46) holds. Since the unstable manifold in  $P_2$  is one-dimensional, we have to choose between two possible directions. The numerical experiment gave us the correct direction. Once we choose  $r$  and compute  $k$ , we have the set  $\kappa$  as given in (42). We should compute the evolution of all points in  $\kappa$  and its intersection with the hyperplane  $v_1 = 4$ . This would require a very long computer time, but since two solutions of (38) cannot intersect, then it is enough to compute the evolution of the points in the boundary of  $\kappa$ , provided we can prove that the trajectories of all points in the interior of  $\kappa$  also reach the hyperplane  $v_1 = 4$ . This can be checked by the following lemma.

LEMMA 29. *Set*

$$(49) \quad \kappa' := \kappa'_{\hat{y}} := \hat{y} + \left\{ \tilde{y} \in S_0 : |\tilde{y}| = \frac{k}{\gamma} r^2 \right\}.$$

Assume that the trajectories of all points in  $\kappa'$  intersect the hyperplane  $v_1 = 4$  and do not intersect the hyperplane  $(2 - n)v_1 - v_2 + 4(n - 2) = 0$ . Let  $\hat{\kappa}$  be the intersection of all such trajectories with  $v_1 = 4$ .

Then the trajectories of all points in  $\kappa$  also intersect the hyperplane  $v_1 = 4$ , and the intersection takes place in the region bounded by  $\hat{\kappa}$ .

*Proof.* Since  $v'_1 = (2 - n)v_1 - v_2 + 4(n - 2)$ , then  $v'_1$  is positive and bounded away from zero for all points of the trajectories starting from  $\kappa'$ . Then, by the uniqueness and continuous dependence on the initial condition of the Cauchy problem, it follows that the union  $\tau(\kappa')$  of such trajectories is a “tube” in  $\mathbb{R}^4$  and the trajectories of all points in  $\kappa \setminus \kappa'$  cannot exit  $\tau(\kappa')$ . Then  $v'_1$  is also positive and bounded away from zero for all points starting in  $\kappa$ , and the trajectory of every point in  $\kappa$  reaches  $v_1 = 4$  in a finite time.  $\square$

Our strategy is as follows: We compute the intersection of the flow starting from all points in  $\kappa'$  with the hyperplane  $v_1 = 4$ . If all the trajectories intersect the hyperplane, we have a proof that the singular solution exists; furthermore the envelope in the  $v_4$ -direction of all intersections yields the desired  $\lambda$ -interval. Note that the set  $\kappa'$  is the image of  $S^2$  through an invertible affine map, and therefore we need an efficient discretization of a sphere.

LEMMA 30. For all  $n = 5, \dots, 16$ , let  $r = .001$ , let  $k$  be as in (48), and, let  $\kappa' = \kappa'_y$  as in (49). For a suitable choice of the direction  $\hat{y}$  in the one-dimensional unstable manifold  $U_0$ , the following conclusions hold:

1. The flow starting in  $\kappa'$  intersects the hyperplane  $v_1 = 4$ .
2. The absolute value of the first coordinate of the intersection point is in the interval set  $[\lambda_\sigma^{\min}, \lambda_\sigma^{\max}]$  defined in Table 1.
3. The flow starting in  $\kappa'$  and ending on the hyperplane  $v_1 = 4$  does not intersect the hyperplane  $(2 - n)v_1 - v_2 + 4(n - 2) = 0$ .

The proof is by computer assistance, as described in section 9.

In order to compute a rigorous lower bound for  $\lambda^*$ , we consider the trajectories of points in the unstable manifold of  $P_1$  and compute the intersection with the hyperplane  $v_1 = 4$ . Since the manifold is two-dimensional, we have to decide the direction to follow: we use the numerical results presented above to compute the direction that gives the highest possible value for  $\lambda$ . We define  $\kappa$  as above, and we wish to prove that all trajectories starting from  $\kappa$  intersect the hyperplane  $v_1 = 4$ . We also need to estimate the location of such intersections. It would save some computer time to restrict the computation to the boundary of  $\kappa$  as in the proof of Lemma 30, but we cannot proceed as in Lemma 29 because  $P_1$  lies on the hyperplane  $v_1 = 4$  and therefore  $v_1$  cannot be monotone. Furthermore, since the unstable manifold has now dimension 2, we do not have the topological argument (the tube) used before. On the other hand, in this case we only have to consider a region which is the affine image of a disk; therefore it is feasible to compute the trajectory for all point in the disk.

LEMMA 31. For all  $n = 5, \dots, 10$ , let  $r = .001$  if  $n \leq 9$  and  $r = .0001$  if  $n = 10$ ; let  $\hat{y} = P_1 + r(e_1 \sin \vartheta_n + e_2 \cos \vartheta_n)$ , where  $e_1$  and  $e_2$  are the eigenvectors of  $A_1$  with unit norm and positive first component corresponding, respectively, to the eigenvalues 2 and 4 and  $\vartheta_5 = 6.2829856$ ,  $\vartheta_6 = 6.28298854$ ,  $\vartheta_7 = 6.2829901$ ,  $\vartheta_8 = 6.2829918$ ,  $\vartheta_9 = 6.2829914$ ,  $\vartheta_{10} = 6.28316589$ ; let  $k$  be as in (48) and let  $\kappa$  be as in (42).

1. The flow starting at all points of  $\kappa$  intersects the hyperplane  $v_1 = 4$ .
2. The absolute value of the first coordinate of the intersection point is larger than the  $\lambda_{\min}^*$  displayed in Table 1.

We point out that this statement shows only that there exists a regular solution

for some value of  $\lambda$  obtained as the intersection of a one-dimensional submanifold of the unstable manifold with the hyperplane  $v_1 = 4$ . Since we cannot exclude that there exists a solution for a larger value of  $\lambda$ , we only have a lower bound for  $\lambda^*$ .

The proof of Theorem 7 follows by Lemmas 27–31.

**7. Proof of Theorem 12.** In this section we use both the PDE notation  $\Delta^2$  and the ODE notation with primes denoting differentiation (with respect to  $r$  or  $s$ , depending on the context).

We assume that  $U_\sigma$  is any radial weakly singular solution of  $(P_{\lambda_\sigma})$  with

$$(50) \quad \lambda_\sigma > 8(n-2)(n-4).$$

In particular, we deal with those solutions obtained in Theorem 7; see also Table 1. Then, by Theorem 6(ii), we know that

$$U_\sigma(r) = -4 \log r + o(|\log r|) \quad \text{as } r \rightarrow 0.$$

Therefore, we define the function

$$W(r) := U_\sigma(r) + 4 \log r$$

and study its behavior. After some calculations, we find that it weakly solves the equation

$$(51) \quad \begin{cases} \Delta^2 W = \frac{1}{|x|^4} [\lambda_\sigma e^W - 8(n-2)(n-4)] & \text{in } B, \\ W = 0 & \text{on } \partial B, \\ \frac{\partial W}{\partial \mathbf{n}} = 4 & \text{on } \partial B. \end{cases}$$

The proof of Theorem 12 follows from the next two lemmas and Proposition 34 at the end of this section.

LEMMA 32. *Assume (50) and assume that  $W \in C^4(0, 1]$  weakly solves (51) ( $W = W(r)$ ); then*

$$(52) \quad \lim_{r \rightarrow 0} W(r) = \log \frac{8(n-2)(n-4)}{\lambda_\sigma} = W_0 < 0.$$

Moreover, at least one of the two following facts holds true:

(i) *The function  $W(r) - W_0$  changes sign infinitely many times in any neighborhood of  $r = 0$ .*

(ii)  *$W(r) \geq \max[W_0, 2r^2 - 2]$  for all  $r \in (0, 1]$ .*

*Proof.* The negativity of  $W_0$  follows from (50), while (52) is a consequence of Theorem 6.

Assume that case (i) in the statement does not occur; we first claim that

$$(53) \quad W(r) \geq W_0 \quad \text{for all } r \in (0, 1].$$

For contradiction, assume that (53) does not hold; then there exists  $\bar{R} \in (0, 1)$  such that  $W(\bar{R}) < W_0$  and two cases may occur, as follows.

*First case.* There exists  $R \in (0, 1)$  such that  $W'(R) = 0$  and  $W_0 \leq W(r) < W(R)$  for all  $r \in (0, R)$ . In this case, let  $H(r) = W(r) - W(R)$  so that  $H(r) < 0$  for all  $r \in (0, R)$ ; on the other hand,  $H$  weakly solves the problem

$$\begin{cases} \Delta^2 H = \Delta^2 W \geq 0 & \text{in } B_R, \\ H = \frac{\partial H}{\partial \mathbf{n}} = 0 & \text{on } \partial B_R, \end{cases}$$

so that by Lemma 16, one gets  $H(r) \geq 0$  for all  $r \in (0, R)$ , a contradiction.

*Second case.* There exists  $R \in (0, 1)$  such that  $W'(R) = 0$  and  $W_0 \geq W(r) > W(R)$  for all  $r \in (0, R)$ . In this case,  $H(r) = W(r) - W(R)$  satisfies both  $H(r) > 0$  for all  $r \in (0, R)$  and

$$\begin{cases} \Delta^2 H = \Delta^2 W \leq 0 & \text{in } B_R, \\ H = \frac{\partial H}{\partial \mathbf{n}} = 0 & \text{on } \partial B_R, \end{cases}$$

giving again a contradiction.

We have so proved (53): hence, if we define the function  $\phi(r) = W(r) + 2 - 2r^2$ , we infer that  $\phi = \phi(|x|)$  weakly satisfies

$$\begin{cases} \Delta^2 \phi = \Delta^2 W \geq 0 & \text{in } B, \\ \phi = \frac{\partial \phi}{\partial \mathbf{n}} = 0 & \text{on } \partial B; \end{cases}$$

this yields  $\phi(r) \geq 0$ , namely,  $W(r) \geq 2r^2 - 2$  for all  $r \in (0, 1]$ .

We have so proved that if (i) does not occur, then (ii) holds true, that is, the statement follows.  $\square$

In high dimensions the previous alternative breaks down, and we can describe the behavior of weakly singular solutions.

LEMMA 33. *If  $n \geq 13$ , then case (i) of Lemma 32 cannot occur.*

*Proof.* Let  $W = W(r)$ , let  $W_0$  be as in Lemma 32, and consider the function

$$Z(s) = W(e^s) - W_0, \quad s \in (-\infty, 0).$$

Then, since  $W$  satisfies (51), we deduce that

$$(54) \quad L_4 Z + p(s)Z = 0, \quad s \in (-\infty, 0),$$

where  $L_4 Z = Z'''' + 2(n - 4)Z''' + (n^2 - 10n + 20)Z'' - 2(n - 2)(n - 4)Z'$  and

$$p(s) = -8(n - 2)(n - 4) \frac{e^{Z(s)} - 1}{Z(s)}.$$

Note that  $p(s)$  is well-defined for all  $s < 0$  and that, by (52),  $p(s) \rightarrow -8(n - 2)(n - 4)$  as  $s \rightarrow -\infty$ . In particular, for all  $\varepsilon > 0$  there exists  $s_\varepsilon < 0$  such that

$$(55) \quad p(s) \geq -[8(n - 2)(n - 4) + \varepsilon] \quad \text{for all } s \leq s_\varepsilon.$$

Since  $n \geq 13$ , for sufficiently small  $\varepsilon$ , the linear equation

$$(56) \quad L_4 Z - [8(n - 2)(n - 4) + \varepsilon]Z = 0$$

admits four linearly independent solutions of “exponential type,” namely,  $Z_i(s) = e^{\nu_i s}$  for some  $\nu_i \in \mathbb{R}$  ( $i = 1, \dots, 4$ ); see also the discussion in section 3.1. Hence, (56) is nonoscillatory in  $(-\infty, 0)$  according to the definition in [E]. Therefore, by (55) and [E, Corollary 1], also (54) is nonoscillatory in  $(-\infty, 0)$  and the statement follows.  $\square$

Let us conclude this section with the observation that an explicit form of the weakly singular solution  $U_\sigma$  seems not so easy to be obtained.

PROPOSITION 34. *Assume that the function  $W$  is a solution of (51) as considered in Lemma 32. Then the function  $W = W(r)$  is not analytic in  $r$  close to 0.*

*Proof.* For contradiction, let  $a_k = W^{(2k)}(0)/(2k)!$  and assume that

$$W(r) = \sum_{k=0}^{\infty} a_k r^{2k}$$

is a convergent power series for  $r$  close to 0. Since  $W$  is regular, the right-hand side of the equation in (51) is bounded as  $r \rightarrow 0$ , and we necessarily have

$$(57) \quad a_0 = \log \frac{8(n-2)(n-4)}{\lambda_\sigma}, \quad a_1 = \frac{W''(0)}{2} = 0.$$

Then

$$W^{(k)}(r) = \frac{W^{(4-k)}(0)}{(4-k)!} r^{4-k} + O(r^{5-k}) \quad \text{as } r \rightarrow 0 \quad (k = 1, 2, 3),$$

and hence

$$\frac{n(n+2)}{3} W^{(4)}(0) = \Delta^2 W|_{r=0} = \frac{\lambda_\sigma e^{a_0} W^{(4)}(0)}{24} = \frac{(n-2)(n-4)}{3} W^{(4)}(0),$$

where we have used (51) and (57). This shows that  $W^{(4)}(0) = 0$  and  $a_2 = 0$ .

We now proceed by induction. Assume that for some  $k \geq 2$  we have shown that  $a_1 = \dots = a_k = 0$ ; we claim that  $a_{k+1} = 0$ . Once we show this, we achieve a contradiction and the statement follows. Note that  $\lambda_\sigma e^W - 8(n-2)(n-4) = 8(n-2)(n-4)[e^{W-a_0} - 1]$  and, by induction assumption,

$$\frac{1}{r^4} (e^{W-a_0} - 1) = a_{k+1} r^{2k-2} + O(r^{2k}).$$

Therefore, from (51) we get

$$(58) \quad \left(\frac{d}{dr}\right)^{2k-2} \Delta^2 W|_{r=0} = 8(2k-2)!(n-2)(n-4)a_{k+1}.$$

On the other hand, recalling the radial form of  $\Delta^2$  (see the left-hand side of (4)) and taking into account that (as  $r \rightarrow 0$ )

$$\begin{aligned} W'(r) &\sim \frac{W^{(2k+2)}(0)}{(2k+1)!} r^{2k+1}, & W''(r) &\sim \frac{W^{(2k+2)}(0)}{(2k)!} r^{2k}, \\ W'''(r) &\sim \frac{W^{(2k+2)}(0)}{(2k-1)!} r^{2k-1}, & W^{(4)}(r) &\sim \frac{W^{(2k+2)}(0)}{(2k-2)!} r^{2k-2}, \end{aligned}$$

we also deduce that

$$\left(\frac{d}{dr}\right)^{2k-2} \Delta^2 W|_{r=0} = 2k(2k+2)(n+2k)(n+2k-2) \cdot (2k-2)! a_{k+1}.$$

Combining this with (58), we get

$$a_{k+1} \{2k(2k+2)(n+2k)(n+2k-2) - 8(n-2)(n-4)\} = 0.$$

Since the term in brackets is strictly positive, this yields  $a_{k+1} = 0$ . □

**8. Further results and open problems.** First, we discuss the stability of the linearizations around regular solutions of problem  $(P_\lambda)$ . For this purpose we observe that the minimal solution depends continuously on  $\lambda$ .

PROPOSITION 35. *As before, let  $U_\lambda$  denote the minimal solution of  $(P_\lambda)$ . Then  $[0, \lambda^*) \ni \lambda \mapsto U_\lambda \in C^{4,\alpha}(\bar{B})$  is continuous from the left. Moreover, if  $\lambda_0 \in [0, \lambda^*)$  is such that the first eigenvalue of the linearization  $L_{U_{\lambda_0}} := \Delta^2 - \lambda_0 \exp(U_{\lambda_0})$  is strictly positive, then  $\lambda \mapsto U_\lambda$  is also continuous in  $\lambda = \lambda_0$ .*

*Proof.* Let  $\lambda_k \nearrow \lambda_0$ . Since  $U_{\lambda_k} \leq U_{\lambda_0}$  and since the  $(U_{\lambda_k})_k$  are monotonically increasing, we get  $\tilde{U} := \lim_{k \rightarrow \infty} U_{\lambda_k}$ , first in any  $L^q$ -space, then by elliptic theory in  $W^{4,q}$ , and finally in  $C^{4,\alpha}(\bar{B})$ . Hence,  $\tilde{U}$  also solves  $(P_{\lambda_0})$ , and  $0 < \tilde{U} \leq U_{\lambda_0}$ . We conclude that  $\tilde{U} = U_{\lambda_0}$  by minimality of  $U_{\lambda_0}$ .

The second statement follows from the implicit function theorem and again the monotonicity of  $U_\lambda$  in  $\lambda$ .  $\square$

The next statement extends some results of [CR] to the biharmonic case; see Proposition 2.15 there. In order to show the sign condition of eigenfunctions, we use a decomposition method with respect to pairs of dual cones.

PROPOSITION 36. *Let  $u$  be a regular solution for  $(P_\lambda)$ , where  $\lambda \in (0, \lambda^*]$ . Let the first eigenvalue  $\mu_1$  of the linearization  $L_u := \Delta^2 - \lambda e^u$  under Dirichlet boundary conditions be nonnegative:  $\mu_1 \geq 0$ . Then every eigenfunction of  $L_u \varphi = \mu_1 \varphi$  is of fixed sign. Moreover, if  $v \in C^4(\bar{B})$  solves  $\Delta^2 v \geq \lambda e^v$  in  $B$  and  $v = \frac{\partial v}{\partial \mathbf{n}} = 0$  on  $\partial B$ , then it follows that  $v \geq u$ . Finally, if  $\mu_1 = 0$ , then we even have  $v = u$ .*

*Proof.* In order to show that the first eigenfunction  $\varphi$  of  $L_u$  is of fixed sign, we need to explain a decomposition technique with respect to dual cones, which was found in the abstract setting by Moreau [Mo] and adapted to biharmonic Dirichlet problems in [GG]. As usual we equip  $H_0^2(B)$  with the scalar product

$$(u, w)_{H_0^2} := \int_B \Delta u \Delta w \, dx.$$

Here, let

$$\mathcal{K} = \{u \in H_0^2(B); u \geq 0 \text{ a.e. in } B\},$$

denote the convex closed cone of nonnegative  $H_0^2$ -functions and

$$\mathcal{K}' = \left\{ u \in H_0^2(B); \text{ for all } w \in \mathcal{K} : (u, w)_{H_0^2} \leq 0 \right\}$$

its dual cone in  $H_0^2$  of weak subsolutions of the clamped plate equation. By Lemma 16 we see that  $\mathcal{K}' \subset -\mathcal{K}$ . For any  $w \in \mathcal{K}'$  we even have that either  $w \equiv 0$  or  $w < 0$  in  $B$ .

Assume now by contradiction that  $\varphi$  is not of fixed sign. Then, according to [Mo], we may decompose

$$\varphi = \varphi_1 + \varphi_2$$

with  $\varphi_1 \in \mathcal{K}$ ,  $\varphi_2 \in \mathcal{K}'$ , and  $\varphi_1 \perp \varphi_2$  in  $H_0^2(B)$ . By assumption we have that  $\varphi_1 \geq 0$ ,  $\varphi_1 \not\equiv 0$ , and  $\varphi_2 < 0$ . But then

$$\begin{aligned} 0 \leq \mu_1 &= \inf_{w \in H_0^2(B) \setminus \{0\}} \frac{\int_B \left( (\Delta w)^2 - \lambda \exp(u) w^2 \right) dx}{\int_B w^2 dx} \\ &\leq \frac{\int_B \left( (\Delta(\varphi_1 - \varphi_2))^2 - \lambda \exp(u) (\varphi_1 - \varphi_2)^2 \right) dx}{\int_B (\varphi_1 - \varphi_2)^2 dx} \end{aligned}$$

$$\begin{aligned} &< \frac{\int_B \left( (\Delta(\varphi_1 + \varphi_2))^2 - \lambda \exp(u)(\varphi_1 + \varphi_2)^2 \right) dx}{\int_B (\varphi_1 + \varphi_2)^2 dx} \\ &= \frac{\int_B \left( (\Delta\varphi)^2 - \lambda \exp(u)\varphi^2 \right) dx}{\int_B \varphi^2 dx} = \mu_1, \end{aligned}$$

a contradiction. Hence,  $\varphi$  is of fixed sign, say  $\varphi \geq 0$ , and in a second step we may conclude from the equation and the strict positivity of the biharmonic Green function (in the ball) that  $\varphi > 0$ .

We consider now  $u$  and  $v$  as in the statement. For  $\tau \in [0, 1]$  we look at

$$(59) \quad \begin{aligned} &\Delta^2(u + \tau(v - u)) - \lambda \exp(u + \tau(v - u)) \\ &\geq \Delta^2(u + \tau(v - u)) - \lambda(\tau \exp(v) + (1 - \tau) \exp(u)) \geq 0. \end{aligned}$$

Since (59) equals 0 for  $\tau = 0$ , its first derivative at  $\tau = 0$  must be nonnegative:

$$(60) \quad \Delta^2(v - u) - \lambda e^u(v - u) =: f \geq 0.$$

If  $\mu_1 > 0$ , a decomposition trick as above applied to the functional  $w \mapsto \int_B ((\Delta w)^2 - \lambda e^u w^2 - f w) dx$  shows that  $v \geq u$ .

If  $\mu_1 = 0$ , we test (60) with the positive first eigenfunction  $\varphi$  and get

$$\Delta^2(v - u) - \lambda e^u(v - u) = 0.$$

That means that also the first derivative of (59) with respect to  $\tau = 0$  vanishes, so that the second derivative needs to be nonnegative:

$$-\lambda e^u (v - u)^2 \geq 0.$$

But this immediately yields  $v = u$ . □

Concerning the stability behavior of the linearizations around regular solutions, we have the following.

**PROPOSITION 37.** *Let  $\lambda > 0$ , let  $u$  be a regular solution of  $(P_\lambda)$ , let  $L_u = \Delta^2 - \lambda e^u$  be the linearized operator at  $u$ , and let  $\mu_1 = \mu_1(L_u)$  be the smallest eigenvalue of  $L_u$ ; then*

- (i) *if  $\lambda < \lambda^*$  and  $u$  is the minimal solution, then  $\mu_1 > 0$ ;*
- (ii) *if  $\lambda < \lambda^*$  and  $u$  is not the minimal solution, then  $\mu_1 < 0$ ;*
- (iii) *if  $\lambda = \lambda^*$  and the extremal solution  $u = U_*$  is regular, then  $\mu_1 = 0$ .*

*Finally, if  $U_\lambda$  denotes the minimal (regular) solution of  $(P_\lambda)$  and  $\mu_1(\lambda) = \mu_1(L_{U_\lambda})$ , then the map  $\lambda \mapsto \mu_1(\lambda)$  is decreasing.*

*Proof.* (i) The monotonicity of  $\mu_1(\lambda)$  follows immediately from the variational characterization

$$\mu_1(\lambda) = \inf_{w \in H_0^2(B) \setminus \{0\}} \frac{\int_B (\Delta w)^2 dx - \int_B \exp(U_\lambda) w^2 dx}{\int_B w^2 dx}$$

and from the monotonicity of  $U_\lambda$  with respect to  $\lambda$ . By Proposition 35 we see that the function  $\lambda \mapsto \mu_1(\lambda)$  is continuous from the left on  $(0, \lambda^*)$  and even on  $(0, \lambda^*]$ , provided the extremal solution  $U_*$  is regular.

Assume by contradiction that there exists a  $\tilde{\lambda} \in (0, \lambda^*)$  with  $\mu_1(\tilde{\lambda}) \leq 0$ . We put

$$\lambda_0 := \sup \{ \lambda \geq 0 : \mu_1(\lambda) > 0 \} \leq \tilde{\lambda} < \lambda^*.$$

According to the mentioned continuity from the left, we have  $\mu_1(\lambda_0) \geq 0$ . If we assume  $\mu_1(\lambda_0) > 0$ , then the second part of Proposition 35 would give  $\mu_1(\lambda) > 0$  also for some  $\lambda > \lambda_0$ , a contradiction. Consequently we have  $\mu_1(\lambda_0) = 0$ . Let  $u = U_{\lambda_0} > 0$  be the corresponding minimal solution:

$$\Delta^2 u = \lambda_0 e^u \text{ in } B, \quad u = \nabla u = 0 \text{ on } \partial B.$$

Consider any  $\lambda \in (\lambda_0, \lambda^*)$  with minimal solution  $v = U_\lambda > 0$ :

$$\Delta^2 v = \lambda e^v \text{ in } B, \quad v = \nabla v = 0 \text{ on } \partial B.$$

Since  $\lambda > \lambda_0$ , Proposition 36 applies and yields  $v = u$  and hence  $\lambda = \lambda_0$ , a contradiction.

(ii) Let  $U_\lambda$  be the minimal solution for  $(P_\lambda)$  so that  $u \geq U_\lambda$ . If the linearization around  $u$  had nonnegative first eigenvalue, then Proposition 36 would also yield  $u \leq U_\lambda$  so that  $u$  and  $U_\lambda$  necessarily coincide, a contradiction.

(iii) Assume that the extremal solution  $u = U_*$  is regular. By continuity, we have  $\mu_1 \geq 0$ . If  $\mu_1 > 0$ , the implicit function theorem would also yield solutions for some  $\lambda > \lambda^*$ . This is a contradiction, so that  $\mu_1 = 0$ .  $\square$

*Open Problem 1.* Does (ii) of Proposition 37 extend to weak solutions  $u$  as formulated in [BV, Theorem 3.1]?

We now turn to the extremal solution  $U_*$ . We first suggest the following open problem.

*Open Problem 2.* Do we have uniqueness of weak solutions for  $(P_{\lambda^*})$ ? By Proposition 37(iii), and arguing as in Lemma 2.6 in [BV], one obtains that if the extremal solution is *regular*, then it is unique even in a weak sense. However, without the regularity assumption on  $U_*$ , the proof seems much more difficult; we refer to [Ma] for the corresponding result related to the second order problem (1). In particular, the proof of a result in the spirit of [Ma, Lemma 2.1] requires a new trick, probably of the same kind as the one we used to prove Lemma 20.

Perhaps, the precise characterization of *all* singular solutions  $U_\sigma$  and the corresponding “singular” parameters  $\lambda_\sigma$  is the most interesting and difficult problem we have to leave open in the present paper.

*Open Problem 3.* Are the singular parameter and the weakly singular solution unique? In order to construct a weakly singular radial solution, according to Theorem 6, one has to follow the unstable branch arising from  $P_2$ . One can do so in two (opposite) exit directions. In one direction we actually find at most (and presumably precisely) one solution by the result of Soranzo [So]: the solution of the PDE has to be strictly decreasing. We emphasize that this result extends to the class of *weakly singular* radial solutions. For the ODE system (8) this means that any “singular” trajectory may intersect the hyperplane  $v_1 = 4$  only once and cannot come back to it. But we do not have a proof that the unstable branch leaving  $P_2$  in the other direction will *not* intersect the hyperplane  $v_1 = 4$  even if numerical experiments suggest so.

Next, we recall that in [GGM] it was shown that for any open bounded domain  $\Omega \subset \mathbb{R}^n$  there exist  $C_1, C_2 > 0$  such that the following improved Hardy inequality holds:

$$(61) \quad \int_{\Omega} |\Delta u|^2 dx \geq \frac{n^2(n-4)^2}{16} \int_{\Omega} \frac{u^2}{|x|^4} dx + C_1 \int_{\Omega} \frac{u^2}{|x|^2} dx + C_2 \int_{\Omega} u^2 dx \quad \text{for all } u \in H_0^2(\Omega).$$

A similar inequality was used in [BV] in order to establish the space dimensions in which the extremal solution for (1) is regular or singular. For  $(P_\lambda)$  this seems more intriguing: it is not clear which is the role of each of the remainder terms in (61). Furthermore, as we have seen in Theorem 12 and Proposition 34, the singular solution is difficult to describe. However, we have a partial result relating Hardy’s inequality with extremal solutions: clearly, this statement is weaker than Corollary 10 if  $n \leq 10$ .

**PROPOSITION 38.** *Let  $\lambda_\sigma$  and  $U_\sigma$  be as in Theorem 7 and assume that  $\lambda_\sigma = \lambda^*$ . Then if  $n \leq 12$ , case (ii) in Theorem 12 cannot occur.*

*Proof.* By Proposition 37(i), by Theorem 3(ii)–(iii), and by using the notation of Theorem 12, we infer that

$$(62) \quad \int_B |\Delta\phi|^2 \geq \lambda^* \int_B e^{U_*} \phi^2 = \lambda^* \int_B \frac{e^W}{|x|^4} \phi^2 \quad \text{for all } \phi \in H_0^2(B).$$

For contradiction, if (ii) in Theorem 12 holds, then

$$\lambda^* \int_B \frac{e^W}{|x|^4} \phi^2 \geq 8(n-2)(n-4) \int_B \frac{\phi^2}{|x|^4} \quad \text{for all } \phi \in H_0^2(B).$$

Since  $8(n-2)(n-4) > \frac{n^2(n-4)^2}{16}$  whenever  $n \leq 12$ , the last inequality, together with (62), would improve the best constant in Hardy’s inequality, a contradiction.  $\square$

Proposition 38 and Corollary 10 suggest the following question and conjecture.

*Open Problem 4.* Which are all the space dimensions  $n \geq 5$  for which  $\lambda_\sigma < \lambda^*$ ? We conjecture that the answer is  $n \leq 12$ . In view of Corollary 10 we know that among these dimensions  $n$ , there are at least  $5 \leq n \leq 10$ . Moreover, Theorem 12 and Proposition 38 prove “half” of this conjecture when  $n = 11, 12$ . Maybe the proof relies on the interpretation of the two remainder terms in (61).

*Open Problem 5.* Show that any radial singular solution is also weakly singular, according to Definition 5. In particular, this would strengthen the statement of Theorem 6.

If the previous three open problems could be solved in the affirmative, then we could also conclude that the extremal solution  $U_*$  is singular if and only if  $n \geq 13$ .

We conclude this paper with some further problems. The next one is not yet completely solved even in the second order case.

*Open Problem 6.* Do there exist singular *nonradial* solutions to  $(P_\lambda)$  for some  $\lambda > 0$ ? We conjecture that the answer is positive; see also Problem 7 in [BV].

Figure 2 displays the numerically computed value of  $-v_4$  of the intersection of a portion of the unstable manifold of  $P_1$  with the hyperplane  $v_1 = 4$  in the case  $n = 5$ .

More precisely,  $-v_4$  is displayed as a function of  $x := -\log(-\vartheta)$ . One may observe the estimated value of  $\lambda^*$  as the maximum value reached by  $-v_4$ ; furthermore, as  $\vartheta \rightarrow 0^-$  the value of  $-v_4$  appears to asymptotically reach  $\lambda_\sigma$  oscillating around it. This leads us to the following problem.

*Open Problem 7.* Assume  $n \leq 12$ . Prove that for every  $N \in \mathbb{N}$  there exists  $\varepsilon = \varepsilon(N) > 0$  such that for  $\lambda \in [\lambda_\sigma - \varepsilon, \lambda_\sigma + \varepsilon]$  there exist at least  $N$  distinct regular radial solutions. For the second order problem the same statement holds true; see [GPP, Theorem 15].

*Open Problem 8.* How can one proceed in arbitrary smooth domains where it is known that comparison principles like Lemma 16 become false? How can one construct and *characterize* the *minimal* solution? Does one have similar bifurcation diagrams, where the solutions, however, can no longer be expected to be positive




FIG. 2.

everywhere? Or does the lack of comparison principles lead to a completely different behavior, at least in geometrically very complicated domains?

**9. Appendix. Computation techniques.** We describe here the algorithm used in the computer assisted proofs. In order to prove Theorem 7 we need a rigorous estimate of the intersection of a branch of the unstable manifold with the hyperplane  $v_1 = 4$ . Since we do not know the exact location of any point of the manifold, except for the stationary point, we compute the trajectory of the whole set  $\kappa'$  as described in section 6. Since no analytical solution of the equation is available, we estimate the trajectories of all points of the set and compute the intersections with the hyperplane  $v_1 = 4$  with rigorous error bounds. In order to compute the image of an infinite set of points, we partition it into boxes with small enough sides, which we call *interval sets*, and we compute their trajectories using interval arithmetics. More precisely, we start with a Taylor approximation of order 10; i.e., we estimate the trajectory of an interval by using the Taylor expansion of order 10 and estimate the error by the Lagrange remainder. If  $h$  is the time step, we compute a rough but rigorous enclosure  $D$  of the trajectory at times  $[0, h]$ , which is an interval set  $D$  such that the solution of the equation lies in  $D$  for all times between 0 and  $h$ . By Lagrange theorem we estimate the error we make neglecting the remaining terms of the Taylor expansion by computing  $x^{(11)}(D) \frac{h^{11}}{11!}$ . We compute  $x^{(11)}(D)$  (which is an interval enclosing all possible values assumed by the 11th derivative of the trajectory, therefore enclosing the Lagrange remainder) using a recursive algorithm for the time derivatives of the solutions (see section I.8 in [HNW]). We point out that it takes a finite amount of  $s$ -time to go from any point in the set  $\kappa'$  (in Lemma 30) or  $\kappa$  (in Lemma 31) to the hyperplane  $v_1 = 4$ . The actual number and size of the intervals that we used as a partition of the sets  $\kappa'$  and  $\kappa$  can be read directly from the *Mathematica* notebook, together with the time step we used for the integration. We feel that it is pointless to display here the long list of numbers which represents such partitions, but since such a list is an essential part of the proof, we make it available in the *Mathematica* notebook.

The interval arithmetics algorithms address the problem of computing the trajectory of an interval and of keeping track of the errors in an elegant and rigorous way, but they introduce another problem. Indeed, even in the simplest dynamical system, the procedure described above leads to a very rough estimate of trajectories, due to the *wrapping effect* which makes the bounds on the error grow exponentially fast. The wrapping effect is one of the main problems one faces when trying to do

rigorous numerics for ODEs.

We describe it with one example: Consider a square centered at the origin  $x = [-\delta, \delta]^2$  and the matrix that represents the rotation in  $\mathbb{R}^2$  by an angle  $\alpha$ ,

$$R(\alpha) = \begin{bmatrix} \cos(\alpha) & -\sin(\alpha) \\ \sin(\alpha) & \cos(\alpha) \end{bmatrix}.$$

Assume for simplicity that  $0 < |\alpha| < \pi/2$ . If we apply  $R_\alpha$  to  $x$  and wish to represent the result by another interval (i.e., another rectangle with sides parallel to the coordinate axes), we see that we need  $(\cos(\alpha) + |\sin(\alpha)|)[- \delta, \delta]^2$ ; therefore, although  $R_\alpha$  is an isometry, its computer realization has a growth factor  $\cos(\alpha) + |\sin(\alpha)| > 1$ . When solving the system of equations of the harmonic oscillator

$$(63) \quad \dot{x} = -y, \quad \dot{y} = x,$$

the  $2\pi$ -shift along the trajectory is an identity map, but when we compute it numerically in interval arithmetics, say with time step  $h = 2\pi/N$ , we have to compose  $N$  times the map induced by  $R(h)$ . An easy computation shows this computation yields a growth factor  $e^{2\pi} \approx 535$  as  $h \rightarrow 0$ .

We substantially reduce the wrapping effect by using the Lohner algorithm. A complete description of interval arithmetics and of the Lohner algorithm is beyond the scope of this paper; we refer to section 6 in [AZ] and the references cited therein for an exhaustive treatment of the topic. More specifically, see [MZ] concerning interval arithmetics and [Lo] for the Lohner algorithm. For the purpose of this description it suffices to consider the Lohner algorithm as a finite number of interval arithmetic operations based on the Taylor expansion which, given (8), an (interval) initial condition  $V_0 \subset \mathbb{R}^4$ , and a time step  $h$ , returns an interval  $V_1 \subset \mathbb{R}^4$  such that for all points  $v_0 \in V_0$  the solution  $v(s)$  of the Cauchy problem with initial condition  $v(0) = v_0$  satisfies  $v(h) \in V_1$ . In other words, the Lohner algorithm provides a rigorous enclosure of the solution at time  $h$  of a given Cauchy problem by performing a finite number of operations. The fact that the operations involved are in finite number and purely arithmetical (they are basically sums and multiplications, which can be performed with computer representable numbers with rigorous control on the round-off) makes it suitable for implementation with a computer.

We must determine the intersection of the trajectory with the hyperplane  $v_1 = 4$ . Since we are computing the trajectory of an interval, it takes a finite (nonzero) amount of “time” to cross the hyperplane; therefore we necessarily introduce another error when estimating the intersection point and have to give a rigorous bound for this error as well. We proceed as follows. We numerically compute the time  $s_1$  required for the flow to reach the intersection. We compute with the Lohner algorithm the solution  $V_1$  of the problem at time  $s_1$ . We check if the first component  $(V_1)_1$  of  $V_1$  is contained in  $(-\infty, 4]$ . If  $(V_1)_1 \subset (-\infty, 4]$ , then no points in  $V_1$  have crossed the hyperplane. If  $(V_1)_1 \not\subset (-\infty, 4]$ , we choose (arbitrarily) a smaller value of  $s_1$  and repeat the step. Then we roughly compute the time  $s_2$  required for the set  $V_1$  to cross the hyperplane. With the Lohner algorithm we compute the solution  $V_2$  of the problem at time  $s_2$ . We check that all points in  $V_2$  have crossed the hyperplane, i.e.,  $(V_2)_1 \subset [4, +\infty)$ . If not, we choose a larger value for  $s_2$  and repeat the step. We are interested only in the value of the fourth component of the solution: since at all points of our interest  $v'_4 < 0$  (because  $v'_4 = v_1 v_4$ ), it suffices to compute the hull of the interval value of  $v_4$  before and after the crossing of the hyperplane. We now have a rigorous proof that the intersection takes place at some  $v_4 \in [\min(V_2)_4, \max(V_1)_4]$ , and this last interval

(with the left bound rounded down and the right bound rounded up) is the value we display in Table 1. For the  $\lambda^*$  computation we display only  $\max(V_1)_4$  rounded down, since the other side of the interval does not have any meaning.

In order to check the third statement in Lemma 30, it is not enough to check that the evolution of all points is in  $A$  as defined in subsection 6.2. Indeed, if the time step is large, it may happen that some trajectory leaves  $A$  and reenters it in a single integration step. We have therefore to check at every time step that the whole rough enclosure  $D$  as defined above is in  $A$  and that the part of the set  $A$  which is contained in the flow tube has a trivial topology, i.e., it does not have holes. The round-off errors are taken care directly by suitable C++ procedures. Such errors may vary by changing computers and/or operating systems, but since they are usually very small when compared to the wrapping effect, we expect that the proofs can be easily reproduced on any recent computer obtaining very similar bounds.

To perform the proofs, we implemented a version of the whole algorithms in a combination of *Mathematica* 4.0 and C++ (gcc version 2.95.1) under the Linux operating system. More precisely, *Mathematica* was used to handle all the data and to perform a few algorithms which are less demanding for the CPU, but more complicated to implement. Furthermore *Mathematica* was used to make all numerical experiments and to draw the pictures. On the other hand C++ was used for the heavy interval arithmetic computations, where it offered much higher speed and more controllable accuracy. The connection between the two languages was obtained by MathLink. The verification of the whole proof takes a few days of CPU time on a machine equipped with an Athlon XP1700 processor. The computer programs which are part of the proofs can be obtained from the authors upon request, while the interval algorithms are provided by [CAPD].

**Acknowledgments.** We are grateful to Anna Dall'Acqua (TU Delft) and Elvise Berchio (Università di Torino) for their careful reading and for helpful remarks. We are also grateful to the referees for their careful review of the paper and for some very useful suggestions on the presentation.

#### REFERENCES

- [ADN] S. AGMON, A. DOUGLIS, AND L. NIRENBERG, *Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions. I*, Comm. Pure Appl. Math., 12 (1959), pp. 623–727.
- [A1] H. AMANN, *Fixed point equations and nonlinear eigenvalue problems in ordered Banach spaces*, SIAM Rev., 18 (1976), pp. 620–709.
- [A2] H. AMANN, *Ordinary Differential Equations. An Introduction to Nonlinear Analysis*, de Gruyter, Berlin, 1990.
- [AZ] G. ARIOLI AND P. ZGLICZYŃSKI, *Symbolic dynamics for the Hénon–Heiles Hamiltonian on the critical level*, J. Differential Equations, 171 (2001), pp. 173–202.
- [BE] J. BEBERNES AND D. EBERLY, *Mathematical Problems from Combustion Theory*, Appl. Math. Sci. 83, Springer, New York, 1989.
- [B] T. BOGGIO, *Sulle funzioni di Green d'ordine  $m$* , Rend. Circ. Mat. Palermo, 20 (1905), pp. 97–135.
- [BCMR] H. BREZIS, T. CAZENAVE, Y. MARTEL, AND A. RAMIANDRISOA, *Blow up for  $u_t - \Delta u = g(u)$  revisited*, Adv. Differential Equations, 1 (1996), pp. 73–90.
- [BV] H. BREZIS AND J. L. VAZQUEZ, *Blow-up solutions of some nonlinear elliptic problems*, Rev. Mat. Univ. Complut. Madrid, 10 (1997), pp. 443–468.
- [C] S. CHANDRASEKHAR, *An Introduction to the Study of Stellar Structure*, Dover, New York, 1967.
- [CAPD] *Computer Assisted Proofs in Dynamics*, <http://capd.wsb-nlu.edu.pl>.

- [CR] M. C. CRANDALL AND P. H. RABINOWITZ, *Some continuation and variational methods for positive solutions of nonlinear elliptic eigenvalue problems*, Arch. Ration. Mech. Anal., 58 (1975), pp. 207–218.
- [E] U. ELIAS, *Nonoscillation and eventual disconjugacy*, Proc. Amer. Math. Soc., 66 (1977), pp. 269–275.
- [GMP] T. GALLOUET, F. MIGNOT, AND J. P. PUEL, *Quelques résultats sur le problème  $-\Delta u = \lambda e^u$* , C. R. Acad. Sci. Paris Sér. I Math., 307 (1988), pp. 289–292.
- [GPP] J. GARCÍA AZORERO, I. PERAL ALONSO, AND J. P. PUEL, *Quasilinear problems with exponential growth in the reaction term*, Nonlinear Anal., 22 (1994), pp. 481–498.
- [GG] F. GAZZOLA AND H.-CH. GRUNAU, *Critical dimensions and higher order Sobolev inequalities with remainder terms*, NoDEA Nonlinear Differential Equations Appl., 8 (2001), pp. 35–44.
- [GGM] F. GAZZOLA, H.-CH. GRUNAU, AND E. MITIDIERI, *Hardy inequalities with optimal constants and remainder terms*, Trans. Amer. Math. Soc., 356 (2004), pp. 2149–2168.
- [G] I. M. GEL'FAND, *Some problems in the theory of quasilinear equations*, Amer. Math. Soc. Transl. (2), 29 (1963), pp. 295–381; translated from the Russian, Uspekhi Mat. Nauk, 14 (1959), pp. 87–158.
- [GNN] B. GIDAS, W. M. NI, AND L. NIRENBERG, *Symmetry and related properties via the maximum principle*, Comm. Math. Phys., 68 (1979), pp. 209–243.
- [GS] H.-CH. GRUNAU AND G. SWEERS, *Positivity properties of elliptic boundary value problems of higher order*, Nonlinear Anal., 30 (1997), pp. 5251–5258.
- [HNW] E. HAIRER, S. P. NØRSETT, AND G. WANNER, *Solving Ordinary Differential Equations I: Nonstiff problems*, 2nd ed., Springer, New York, 2000.
- [J] J. JACOBSEN, *A Liouville–Gelfand equation for  $k$ -Hessian operators*, Rocky Mountain J. Math., 34 (2004), pp. 665–684.
- [JS] J. JACOBSEN AND K. SCHMITT, *The Liouville–Bratu–Gelfand problem for radial operators*, J. Differential Equations, 184 (2002), pp. 283–298.
- [JL] D. JOSEPH AND T. S. LUNDGREN, *Quasilinear Dirichlet problems driven by positive sources*, Arch. Ration. Mech. Anal., 49 (1973), pp. 241–269.
- [Li] P.-L. LIONS, *On the existence of positive solutions of semilinear elliptic equations*, SIAM Rev., 24 (1982), pp. 441–467.
- [Lo] R. J. LOHNER, *Computation of guaranteed enclosures for the solutions of ordinary initial and boundary value problems*, in Computational Ordinary Differential Equations, J. R. Cash and I. Gladwell, eds., Clarendon Press, Oxford, 1992, pp. 425–435.
- [Ma] Y. MARTEL, *Uniqueness of weak extremal solutions for nonlinear elliptic problems*, Houston J. Math., 23 (1997), pp. 161–168.
- [MP1] F. MIGNOT AND J. P. PUEL, *Sur une classe de problèmes non linéaires avec non-linéarité positive, croissante, convexe*, Comm. Partial Differential Equations, 5 (1980), pp. 791–836.
- [MP2] F. MIGNOT AND J. P. PUEL, *Solution radiale singulière de  $-\Delta u = \lambda e^u$* , C. R. Acad. Sci. Paris Sér. I Math., 307 (1988), pp. 379–382.
- [Mo] J. J. MOREAU, *Décomposition orthogonale d'un espace hilbertien selon deux cônes mutuellement polaires*, C. R. Acad. Sci. Paris, 255 (1962), pp. 238–240.
- [MZ] M. MROZEK AND P. ZGLICZYŃSKI, *Set arithmetic and the enclosing problem in dynamics*, Ann. Polon. Math., 74 (2000), pp. 237–259.
- [PT] L. A. PELETIER AND W. C. TROY, *Spatial patterns. Higher order models in physics and mechanics*, Progr. Nonlinear Differential Equations Appl. 45, Birkhäuser, Boston, Boston, MA, 2001.
- [P] S. I. POHOŽAEV, *Eigenfunctions of the equation  $\Delta u + \lambda f(u) = 0$* , Soviet Math. Dokl., 6 (1965), pp. 1408–1411.
- [PS] P. PUCCI AND J. SERRIN, *A general variational identity*, Indiana Univ. Math. J., 35 (1986), pp. 681–703.
- [So] R. SORANZO, *A priori estimates and existence of positive solutions of a superlinear polyharmonic equation*, Dynam. Systems Appl., 3 (1994), pp. 465–487.
- [Sw] G. SWEERS, *No Gidas–Ni–Nirenberg type result for biharmonic problems*, Math. Nachr., 246/247 (2002), pp. 202–206.
- [We] J. WEI, *Asymptotic behavior of a nonlinear fourth order eigenvalue problem*, Comm. Partial Differential Equations, 21 (1996), pp. 1451–1467.
- [Wi] CH. WIENERS, *Numerische Existenzbeweise für Schwache Lösungen Nichtlinearer Elliptischer Randwertaufgaben (Numerical proofs of existence of weak solutions of nonlinear elliptic boundary value problems)*, Ph.D. thesis, University of Cologne, 1994.